

A modified game of 'chicken' reveals what happens in the brain during decision-making

December 2 2020, by Michele W. Berger



Credit: Pixabay/CC0 Public Domain

Better understanding what happens in the brain when people cooperate and make social decisions has the potential to help boost such behaviors. It's been a research focus for the University of Pennsylvania's Platt Labs



for some time.

"The social brain network contains a number of different areas critical to managing interactions and connecting with others," says Penn Integrates Knowledge Professor Michael Platt. "One is more emotional, critical for empathy and learning from others. That's the anterior cingulate gyrus. The other is more cognitive, and for humans that lives in the temporoparietal junction."

In nonhuman primates, the anterior cingulate gyrus mirrors what it does in humans, responding to rewards and pain, both for themselves and others. Until recently, scientists believed that nonhuman primates did not have an equivalent to the temporoparietal junction, which activates when humans think about the beliefs, desires, and goals of others. But new research on <u>rhesus macaques</u> from Platt, postdoctoral fellow Wei Song Ong, and Duke University postdoctoral fellow Seth Madlon-Kay confirms that an area called the middle superior temporal sulcus (mSTS) actually plays a similar role.

What's more, analyzing the firing rates of neurons in those two sections of the brain as the macaques engage in a task together—one in which the reward for each participant depends, in part, on how the two cooperate—reveals a high level of strategic decision-making, findings Platt and colleagues published in the journal *Nature Neuroscience*.

To test the research team's theories about the role of these brain regions, Ong created an experiment akin to a modified game of chicken. In the best-known version, participants face off and move toward each other in actual cars; the one who swerves to avoid the collision is the chicken. "Here, each monkey uses a joystick to control a circle or ring, which we call a car," Platt says. "They have about 10 seconds to decide whether to drive the car straight or turn it to the side."



In either direction, the macaques see tokens proportional to the juice reward they would receive for driving to each spot. "The temptation is to drive forward because there is typically more reward there," he says. "But if they both drive forward, they 'crash," and no one gets a reward." The researchers also placed a white bar behind the side tokens; if the participants coordinated their behavior, turning to that spot at the same time, they could unlock the extra reward.

"It's a way to give them the opportunity to work together," Platt says. "It's never more <u>reward</u> than going forward, but it's an intermediate. It's better than losing."

With the study design set, the Penn team conducted tens of thousands of trials for three conditions: one with two <u>active participants</u>, a second with an active participant and a computer, and third with an active participant and a decoy participant that sat in the opposite seat and shared the rewards but never actually controlled the joystick.

The researchers observed the behavior in each trial to determine, first and foremost, whether the macaques understood the rules. "The initial pass at the data shows they do," says Platt. "They follow the payoffs. They're more or less performing optimally." The researchers then built a computational model of the behavior, progressively adding in components—previous actions, number of tokens available, actions of the partner participant—to explain what was happening. Finally, they used the parameters from the model as variables in trying to understand what information the neurons in each area were encoding.

Taking this all into account, they discovered that for <u>nonhuman primates</u> the anterior cingulate gyrus and mSTS behave similarly to the anterior cingulate gyrus and the <u>temporoparietal junction</u> in humans, specifically as they relate to coordination and high-level decision-making.



"This is a really interesting outcome," Platt says. "It tells us a lot about how sophisticated monkeys are in terms of their social reasoning. It tells us something about the process unfolding in the brain as we interact with another individual and make decisions for or against each other. And it tells us that our ability to cooperate, to work together in societies, emerged from ancestral adaptations that evolved some 20 million years ago."

In the future, Platt says he would like to test the findings on more realistic tasks and situations, to determine whether they hold up in realworld scenarios.

For now, the latest research brings the team nearer to comprehending what happens in the brain during humans' decision-making process. "We are closer to understanding, in a general sense, how this system works," he says, "how differences in the way this system functions in different people might be associated with different types of <u>social decisions</u>."

More information: Wei Song Ong et al. Neuronal correlates of strategic cooperation in monkeys, *Nature Neuroscience* (2020). DOI: 10.1038/s41593-020-00746-9

Provided by University of Pennsylvania

Citation: A modified game of 'chicken' reveals what happens in the brain during decision-making (2020, December 2) retrieved 27 June 2024 from <u>https://medicalxpress.com/news/2020-12-game-chicken-reveals-brain-decision-making.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.