

Genes play a role in common knee injury

December 15 2020



Credit: CC0 Public Domain

It has long been known that the choice of shoe, surface and type of sport can all be contributing factors when someone suffers an anterior cruciate ligament rupture. Researchers at Lund University in Sweden have now observed that genes also play a decisive role.

By analyzing data from the Swedish Twin Register along with data from the Swedish National Patient Register, researchers were able to see how many [identical twins](#) and fraternal twins had been diagnosed with anterior cruciate ligament rupture. A total of 88 414 twins aged 17 or above were involved in the [epidemiological study](#), published in the British Journal of Sport Medicine.

The opportunity to observe injuries among both identical and fraternal twins enables conclusions to be drawn about heredity, explains Karin Magnusson, researcher and expert in [twin studies](#) within the field of orthopedics at Lund University.

"Identical twins have totally identical genes, while fraternal twins, like other siblings, share half of their genes with each other. We can estimate heritability by studying how often cruciate ligament injury occurs in both twins in a set, that is, we compare the 'double' prevalence in identical and [fraternal twins](#). This way we can draw conclusions about the importance of heredity and environment."

"Our results show that genes appear to contribute more than we thought. To put it very simply, we can say that out of all cruciate ligament injuries, 69 percent can be explained by genetics. This should not be interpreted to mean an individual's risk of suffering the injury is 69 percent, rather that it is significant in terms of the wider population. Heredity is easier to understand when compared with other illnesses or conditions. The [genetic risk](#) of suffering from cancer is 33 percent, for example. This means that 33 percent of variation within the population in terms of causes of all cancer cases is due to [genetic variation](#), while the rest is due to other factors—such as environment or lifestyle," explains Magnusson.

The genetic risk was equally high in men and women.

The results may be significant in terms of preventing this type of injury more effectively.

"In the past, genetic factors have not been taken into account when working to prevent anterior cruciate ligament injury, for example among athletes. This study does not provide us with an answer as to whether heredity entails anatomical or physiological conditions that could affect the risk of a cruciate [ligament](#) injury. However, if we know that there are many cases of this [injury](#) within a family, then it is worth being extra cautious and doing more preventive training, such as landing after jumping when playing handball, and so on," says Martin Englund, professor at Lund University, physician at Skåne University Hospital and one of the researchers behind the study.

More information: Karin Magnusson et al. High genetic contribution to anterior cruciate ligament rupture: Heritability ~69%, *British Journal of Sports Medicine* (2020). [DOI: 10.1136/bjsports-2020-102392](https://doi.org/10.1136/bjsports-2020-102392)

Provided by Lund University

Citation: Genes play a role in common knee injury (2020, December 15) retrieved 6 May 2024 from <https://medicalxpress.com/news/2020-12-genes-role-common-knee-injury.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--