

Study shows link between cervical cancer and HIV infection

December 21 2020



Women infected with HIV are six times more likely to develop cervical cancer. This is what Dr. Dominik Stelzle and Dr. Luana Tanaka found out as part of a study by the Center for Global Health at the Technical University of Munich. Credit: Andreas Heddergott / TUM

A research team at the Technical University of Munich (TUM) has

quantified the effects of an infection with the Human Immunodeficiency Virus (HIV) on the development of cervical cancer. Their results show that the risk of developing cervical cancer is six times higher in women who are infected with HIV. Southern and Eastern Africa are particularly affected.

According to WHO statistics, cervical [cancer](#) is the fourth most common type of cancer for women. In 2018 an estimated 570,000 women worldwide were diagnosed with cervical carcinoma, with approximately 311,000 of these women dying.

On the other hand cervical cancer, usually caused by Human Papillomavirus (HPV), is also one of the most successfully preventable and treatable types of cancer, as long as it is detected at an early stage and treated effectively.

Cervical cancer is at the same time the most frequently detected cancer for women who live with HIV, since their immune systems are weakened by the HIV [infection](#).

The TUM School of Medicine's Center for Global Health and the Chair of Epidemiology at the TUM Department of Sport and Health Sciences have now dedicated their efforts to this relevant topic in the publication "Estimates of the Global Burden of Cervical Cancer Associated with HIV", published in the renowned journal *The Lancet Global Health*.

Systematic review and meta-analysis of 24 studies

The lead authors Dr. Dominik Stelzle (Center for Global Health and Chair of Epidemiology) and Dr. Luana Tanaka (Chair of Epidemiology) conducted a [systematic review](#) as well as a meta-analysis of a total of 24 studies from the years 1981 to 2016, in which 236,127 women with HIV from four continents (Africa, North America, Asia and Europe)

participated.

These studies covered a total of 2,138 cervical carcinoma cases. The results were linked with data from UNAIDS on worldwide HIV infection and with data on cervical carcinoma from the International Agency for Research on Cancer (IARC), the WHO's Cancer Research Center, and then evaluated.

"Until now there have only been estimates from countries with high net income levels," says Dr. Stelzle. "That's why we looked at the figures on global incidence of cervical carcinoma in connection with an HIV infection and included estimates for countries with low net incomes. In most parts of the world the numbers are under five percent. In some countries however we're talking about well over 40 percent of cases."

Risk is six times higher for women with HIV

The objective of the study was to calculate the share of women living with HIV among the number of women with cervical cancer. The authors found that 5.8 percent of all new cervical cancer cases worldwide in the year 2018 were diagnosed for women with an HIV infection. This is equivalent to 33,000 cases a year, 85 percent of which occurred in Sub-Saharan Africa.

Furthermore, based on their results the team was able to show that women with HIV have a sixfold higher risk of developing cervical cancer than women without HIV infection.

"The association between cervical [carcinoma](#) and HIV is plausible," says Prof. Andrea S. Winkler, co-director of the Center for Global Health. "Cervical carcinomas are usually caused by infections with Human papillomavirus (HPV), which are sexually transmitted just as HIV is. Based on our results it can be assumed that an infection with HIV

represents a risk factor for an infection with HPV."

Southern and eastern Afrika most affected

The regions most strongly affected are southern and eastern Africa, where respectively 63.8 percent and 27.4 percent of cervical carcinomas were diagnosed in women with an HIV infection.

"With over 75 percent, Eswatini in southern Africa is the country with the highest share of women suffering from cervical cancer in connection with an HIV infection, followed by Lesotho with 69 percent, Botswana with 67 percent, South Africa with 64 percent and Zimbabwe with 52 percent," says Dr. Tanaka.

Based on the results, the TUM authors determined that women with an HIV infection have a significantly higher risk of developing cervical cancer. They also pointed out that this means that HPV vaccinations and early-stage [cervical carcinoma](#) screenings are of particular importance, especially in the African countries south of the Sahara.

"Although [cervical cancer](#) screenings already exist in Africa, until now they have been performed primarily on women with a higher socio-[economic status](#) who could afford the screening," says Prof.

Stefanie Klug, who holds the Chair of Epidemiology at TU Munich.

"The goal has to be to end this dependency on economic means and to succeed in introducing free HPV vaccination for girls and screening for adult [women](#)."

More information: Correction to Lancet Glob Health 2020; published online Nov 16. [doi.org/10.1016/S2214-109X\(20\)30459-9](https://doi.org/10.1016/S2214-109X(20)30459-9), *The Lancet Global Health* (2020). [DOI: 10.1016/S2214-109X\(20\)30509-X](https://doi.org/10.1016/S2214-109X(20)30509-X)

Provided by Technical University Munich

Citation: Study shows link between cervical cancer and HIV infection (2020, December 21)
retrieved 27 April 2024 from

<https://medicalxpress.com/news/2020-12-link-cervical-cancer-hiv-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.