

Oral contraceptive pills protect against ovarian and endometrial cancer

December 17 2020



Credit: CC0 Public Domain

A comprehensive study from Uppsala University involving more than 250,000 women shows that oral contraceptive use protects against ovarian and endometrial cancer. The protective effect remains for

several decades after discontinuing the use. The study is published in the journal *Cancer Research*.

Ovarian and [endometrial cancer](#) are among the most common gynecological cancers, with a lifetime risk of just over 2%. Endometrial cancer is slightly more common but as it has clearer symptoms and is therefore often detected at an early stage, the mortality rate is low. However, [ovarian cancer](#) is among the deadliest cancers, since it is often not detected until it has already spread to other parts of the body.

The first oral contraceptive pill was approved already in the 1960s, and 80% of all women in Western Europe have used [oral contraceptives](#) at some point in their life. Oral contraceptives include estrogen and progestin, which are synthetic forms of the female sex hormones. The estrogen and progestin in oral contraceptives prevent ovulation and thereby protect against pregnancy.

In the current study, the scientists compared the incidence of breast, ovarian and endometrial cancers between women that had used [oral contraceptive pills](#) and never users.

"It was clear that women who had used oral contraceptive pills had a much [lower risk](#) of developing both ovarian and endometrial cancer. Fifteen years after discontinuing with oral contraceptives, the risk was about 50% lower. However, a decreased risk was still detected up to 30-35 years after discontinuation," says Åsa Johansson at the Department of Immunology, Genetics and Pathology, Uppsala University, one of the leading researchers behind the study.

However, oral contraceptive pills have previously been associated with an increased risk of breast cancer.

"Surprisingly, we only found a small increased risk of breast cancer

among oral contraceptive users, and the increased risk disappeared within a few years after discontinuation," says Johansson. "Our results suggest that the [lifetime risk](#) of breast cancer might not differ between ever and never users, even if there is an increased short-term risk."

The results from the current study are important, since oral contraceptive use has commonly been associated with side effects such as deep vein thrombosis and breast cancer.

"In addition to protecting against pregnancy, we have shown that oral contraceptive pills also have other positive effects. Our results can enable women and physicians to make more informed decisions about which women should use oral contraceptive pills," says Therese Johansson, one of the Ph.D. students behind the study.

More information: Karlsson T, Johansson T, Höglund T, Ek W, Johansson Å. (2020) Time-dependent effects of oral contraceptive use on breast, ovarian and endometrial cancers. *Cancer Research* (2020). [DOI: 10.1158/0008-5472.CAN-20-2476](https://doi.org/10.1158/0008-5472.CAN-20-2476)

Provided by Uppsala University

Citation: Oral contraceptive pills protect against ovarian and endometrial cancer (2020, December 17) retrieved 27 April 2024 from <https://medicalxpress.com/news/2020-12-oral-contraceptive-pills-ovarian-endometrial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
