


Racially biased oxygen readings could be putting patients at risk

December 16 2020



Credit: CC0 Public Domain

Thousands of times a day, doctors and nurses use a small device called a pulse oximeter—which fits on the end of a patient's finger to measure the percentage of oxygen in the blood—to help them make vital treatment decisions. However, a new study by Michigan Medicine researchers finds that the pulse oximeter may be less accurate in Black patients than in White patients.

"These results really surprised all of us," says Michael Sjoding, M.D., assistant professor of internal medicine in the division of pulmonary and [critical care medicine](#). As he and his colleagues cared for critically ill patients with COVID-19, a significant percentage of whom were Black, they noted a reoccurring discrepancy between readings from the [pulse oximeter](#) and a measurement from an arterial [blood gas](#), which measures [oxygen saturation](#) directly through a [blood test](#).

Using a data from patients hospitalized at Michigan Medicine and an older dataset of patients hospitalized in multiple intensive care units across the country, the research team compared results from pulse oximeters and arterial blood gas measurements in patients who identified as either Black or White.

"We wanted to know: when the pulse oximeter is measuring between 92 and 96 percent, how often is the oxygen saturation actually less than 88 percent on arterial blood gas?" says Sjoding, adding that a reading of below 88 percent is generally where most physicians start to take notice and intervene.

Black patients were three times as likely to have a significant discrepancy in these readings. Twelve percent of the time, when Black patients had a pulse oximeter reading in this safe range, their actual saturation on an arterial blood gas was below 88%. In [white patients](#), this discrepancy occurred three percent of the time.

"Many studies have examined health disparities between White and Black patients, with most determining Black patients have worse outcomes than White patients. What these studies often lack are reasons to explain these differences," says study co-author Thomas Valley, M.D., Assistant Professor in the Division of Pulmonary and Critical Care Medicine in the Department of Internal Medicine at the University of Michigan. "Our findings may provide one key explanation."

The findings raise a number of important questions about the management of patients in the hospital. Sjoding notes that there has been a departure from using the more invasive arterial blood gas measurements in favor of the simpler, less invasive pulse oximeter to determine whether a patient is getting enough oxygen. Knowing that pulse oximeter readings may be inaccurate will "absolutely affect therapy," he says. "I was caring for critically ill patients in the Intensive Care Unit when I first ran this analysis. The next day I started to rethink all my management strategies."

The findings aren't entirely new, the team reports. A paper published in an anesthesia journal in 2005 also reported that skin pigmentation led to inaccuracy in pulse oximeter readings. "At the time, these findings weren't considered significant enough to change practice," says Sjoding.

"I think we're going to have to be more cautious about whether Black patients are truly getting that amount of oxygen they need. We may need to pay more attention to arterial blood gas results to confirm that the patient is getting enough oxygen."

Sjoding says longer term, device makers may need to make additional changes to ensure that all patients are getting an accurate reading. Inaccurate pulse oximeter readings could have implications for outpatient care as well, as most clinics rely on the measurements to determine treatment needs.

"We're concerned about patients who need supplemental oxygen at home, like patients with emphysema. Those patients can qualify for supplemental [oxygen](#) if their saturation is less than 88 percent," says Sjoding. "There's no reason to think that these findings wouldn't apply to patients in an outpatient clinic."

More information: *New England Journal of Medicine*, [DOI](#):

[10.1056/NEJMc2029240](https://doi.org/10.1056/NEJMc2029240)

Provided by University of Michigan

Citation: Racially biased oxygen readings could be putting patients at risk (2020, December 16)
retrieved 5 May 2024 from

<https://medicalxpress.com/news/2020-12-racially-biased-oxygen-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.