

Vitamin D the clue to more autism spectrum disorder in boys

December 11 2020



Credit: Pixabay/CC0 Public Domain

A deficiency in Vitamin D on the mother's side could explain why autism spectrum disorder is three times more common in boys, say researchers from The University of Queensland.

In their latest study, Professor Darryl Eyles and Dr. Asad Ali from UQ's

Queensland Brain Institute found vitamin D deficiency during pregnancy caused an increase in testosterone in the developing brain of male rats.

"The biological cause of Autism spectrum disorder (ASD) is unknown but we have shown that one of the many risk factors—low vitamin D in mothers—causes an increase in testosterone in the brain of the male fetuses, as well as the maternal blood and [amniotic fluid](#)," Professor Eyles said.

"In addition to its role in calcium absorption, vitamin D is crucial to many developmental processes.

"Our research also showed that in vitamin D-deficient male fetuses, an enzyme which breaks down testosterone was silenced and could be contributing to the presence of high testosterone levels."

Professor Eyles' previous research has shown that vitamin D plays a critical role in [brain development](#) and that that giving vitamin D supplements to mice during pregnancy completely prevented ASD-like traits in their offspring.

Co-author Dr. Ali said that excessive exposure of the developing brain to sex hormones like testosterone was thought to be an underlying cause of ASD, but the reasons remained unclear.

"Vitamin D is involved in pathways controlling many sex hormones," Dr. Ali said.

"When the rat mothers were fed a low vitamin D diet, it caused male fetal brains to have high levels of exposure to testosterone."

Professor Eyles said the study was the first to show that a known risk factor for ASD alters testosterone in both the fetal brain and the

mother's blood—one possible contributor to why ASD is more prevalent in males.

"We have only studied one risk factor for ASD—vitamin D deficiency during development—our next step is to look at other possible [risk factors](#), such as maternal stress and hypoxia—lack of oxygen—and see if they have the same effect," he said.

More information: Asad Amanat Ali et al, Developmental vitamin D deficiency increases foetal exposure to testosterone, *Molecular Autism* (2020). [DOI: 10.1186/s13229-020-00399-2](https://doi.org/10.1186/s13229-020-00399-2)

Provided by University of Queensland

Citation: Vitamin D the clue to more autism spectrum disorder in boys (2020, December 11) retrieved 23 April 2024 from <https://medicalxpress.com/news/2020-12-vitamin-d-clue-autism-spectrum.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.