

# Weekly folic acid boost shows potential to halve birth defects

December 7 2020

---



Credit: Pixabay/CC0 Public Domain

SAHMRI researchers have found the risk of birth defects can be greatly reduced if women significantly increase their folic acid intake.

Neural tube defects (NTDs) are [birth defects](#) like [spina bifida](#) that affect 300,000 babies born annually worldwide. Some of these babies die at birth while others have life-long disabilities.

The study sponsored by Nutrition International with support from the Government of Canada and published in *BMJ Global Health*, has shown taking weekly iron [folic acid supplements](#) (IFA) containing 2.8 mg folic acid can lower the risk of NTDs by up to four times more than the current global standard of just 0.4 mg.

Project leader, Professor Tim Green said during the trial, 70% of women consuming 2.8 mg folic acid per week achieved a red blood cell folate concentration associated with a low NTD risk, compared with just 10% of those taking 0.4 mg.

"A folic acid supplement should be taken not only by those who are planning a pregnancy, but by all women who are sexually active," Prof Green said.

"We know around half of pregnancies are unplanned globally and in low and middle income countries 10 million of these pregnancies occur each year in adolescents alone. By the time these women realize they're pregnant it's too late to get the benefits of IFA."

Researchers predict the number of NTDs worldwide could be halved if the World Health Organization (WHO) acts on the latest findings.

The WHO currently recommends a weekly supplement for all women aged 15-49 living in countries where the rate of anemia is above 20%. This supplement currently contains the appropriate 60mg of iron but just 0.4mg of folic acid; not enough to provide a benefit.

"It's our recommendation that the optimal 2.8mg of [folic acid](#)

formulation be made widely available to women in low and [middle income countries](#) as soon as possible," Prof Green said.

**More information:** Kaitlyn L I Samson et al. Weekly iron–folic acid supplements containing 2.8 mg folic acid are associated with a lower risk of neural tube defects than the current practice of 0.4 mg: a randomized controlled trial in Malaysia, *BMJ Global Health* (2020). [DOI: 10.1136/bmjgh-2020-003897](#)

Provided by BMJ Global Health

Citation: Weekly folic acid boost shows potential to halve birth defects (2020, December 7) retrieved 21 May 2024 from <https://medicalxpress.com/news/2020-12-weekly-folic-acid-boost-potential.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.