

Abnormal hyperactivation in the brain may be an early sign of Alzheimer's

January 21 2021, by Salle de Presse



Credit: Unsplash/CC0 Public Domain

Abnormally hyperactive areas in the brain may help better predict the onset of Alzheimer's disease, according to findings of a research team led by Université de Montreal psychology professor Sylvie Belleville,

scientific Director of the Institut universitaire de gériatrie de Montréal research centre.

Hyperactivation could be an early biomarker of Alzheimer's [disease](#), the researchers say in their study published today in *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, co-authored by Belleville and Nick Corriveau-Lecavalier, a doctoral student she supervises.

Worried about their memory

In their research, the team found hyperactivation in certain [brain areas](#) in people not yet diagnosed with Alzheimer's but who were worried about their memory and who exhibited risk factors for the disease.

The study marks an important milestone in this research area, as the hyperactivation of regions susceptible to the Alzheimer's as shown by functional magnetic resonance imaging (fMRI) was observed in people with no clinical symptoms and before the onset of cognitive impairments detected with standardized tests.

"This study indicates that abnormal activation in these areas may be observed many years before diagnosis," said Belleville.

This finding is crucial to the advancement of knowledge about the disease, she continued.

"Alzheimer's disease is progressive and may emerge in the brain 20 to 30 years before diagnosis. It is therefore very important to pinpoint biomarkers—that is, physical and detectable signs of the disease—and to better understand the initial effects on the brain. Hyperactivation could therefore represent one of the first signs of Alzheimer's disease."

An inverse U-shape

The team observed that, as the disease progresses, neuronal activation follows an inverse U-shape trajectory. Indeed, activation in certain areas of the brain in the early stages of Alzheimer's may significantly increase before the neuronal loss that is caused by the disease leads to a clear decrease in activation.

"This form may characterize the underlying pathological process and help doctors determine the stage of the disease," explained Corriveau-Lecavalier, the study's first author. "When combined with other indicators such as blood work and cognitive tests, this type of neuroimaging investigation could help with possible earlier detection."

For their study, the team used data from the Consortium for the Early Identification of Alzheimer's Disease to study brain activation in [groups of individuals](#) at a high risk of developing Alzheimer's disease who had performed a memory task while being scanned with fMRI. One group consisted of 28 individuals who were concerned about their memory but who did not show cognitive impairments on traditional clinical tests. The other group included 26 individuals with mild cognitive impairments.

The researchers found that the individuals in the first group, or those with memory complaints but who did not show objective cognitive impairments, had abnormally high levels of activation in multiple key regions of the brain affected by Alzheimer's disease. Individuals with mild cognitive impairments, who are considered to be at a more advanced stage of the disease, tended to show decreased activation in these [brain](#) regions.

More information: Nick Corriveau-Lecavalier et al, A quadratic function of activation in individuals at risk of Alzheimer's disease, *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*

(2021). [DOI: 10.1002/dad2.12139](https://doi.org/10.1002/dad2.12139)

Provided by University of Montreal

Citation: Abnormal hyperactivation in the brain may be an early sign of Alzheimer's (2021, January 21) retrieved 2 May 2024 from <https://medicalxpress.com/news/2021-01-abnormal-hyperactivation-brain-early-alzheimer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.