

At-home swabs diagnose infections as accurately as healthcare worker-collected swabs

January 28 2021



Credit: CC0 Public Domain

Self swabs and caregiver swabs are effective at detecting multiple pathogens and are just as accurate as those taken by healthcare workers,

according to a team of Australian researchers. The research appears in the *Journal of Clinical Microbiology*, a publication of the American Society for Microbiology.

"Across the range of pathogens and swab types, there was high agreement between results from self- or caregiver swabs and those performed by a [healthcare worker](#), even when different sites were swabbed (e.g. nasopharyngeal vs. nasal)," said principal investigator Joshua Osowicki, BMedSci, MBBS, FRACP, a Pediatric Infectious Diseases physician in the Murdoch Children's Research Institute's Tropical Diseases research group, Melbourne, Australia.

"We found a pooled sensitivity of 91% and specificity of 98% for self- or caregiver-collected upper airway swabs compared to [[healthcare worker](#)] swabs," the investigators wrote. (Sensitivity refers to the proportion of those who test positive that are correctly identified, and specificity the proportion testing negative that are correctly identified..)

"Findings were similar when only SARS-CoV-2 data were considered," the researchers wrote. "This level of diagnostic performance should reassure clinicians, researchers, and public health officials that diagnostic performance is not necessarily compromised by self- or caregiver swabbing."

Pathogens swabbed in the study include SARS-CoV-2, influenza, group A Streptococcus, Staphylococcus aureus, pharyngeal Chlamydia trachomatis and Neisseria gonorrhoeae, and a panel of respiratory viruses.

The study was a [systematic review](#) and meta-analysis of 20 previous studies comprising swabs of more than 3,500 individuals, comparing self- or caregiver swabs with swabs collected by healthcare workers for diagnosis of a range of upper airway pathogens. "We searched for

previous studies that included both kinds of swabs collected from the same individuals," said Dr. Osowicki. "For each study, we considered the sensitivity and specificity of self- or caregiver swabs compared to swabs by healthcare workers, and the agreement between the two kinds of swabs.

Self- and caregiver swabbing offer a number of advantages over swabbing by healthcare workers, including less likelihood of transmission, reduced expenses, conservation of personal protective equipment, and less discomfort to patients. An online survey of patients with SARS-CoV-2 symptoms found a much higher proportion willing to be tested via a self-collected saliva sample (92%) or a nasal swab (88%) than by a drive-through [healthcare](#) worker [swab](#) collection (71%)," according to the investigators.

When the plans of then [medical student](#) and first author Ciara Harrison to complete an immunology laboratory project were derailed by the pandemic, Dr. Harrison, now a first-year physician, pivoted to this systematic review and meta-analysis.

More information: Ciara Harrison et al, A systematic review and meta-analysis of upper airways swab collection for detection of viral and bacterial pathogens by individuals or caregivers compared to healthcare workers., *Journal of Clinical Microbiology* (2021). [DOI: 10.1128/JCM.02304-20](#)

Provided by American Society for Microbiology

Citation: At-home swabs diagnose infections as accurately as healthcare worker-collected swabs (2021, January 28) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-01-at-home-swabs-infections-accurately-healthcare.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.