

New research highlights dangerous disparities for life-saving cancer screening

January 13 2021

JOURNAL OF THE NATIONAL COMPREHENSIVE CANCER NETWORK



NCCN Guidelines®

- GENETIC/FAMILIAL HIGH-RISK ASSESSMENT: BREAST, OVARIAN, AND PANCREATIC

NCCN Guidelines® Insights

- ACUTE MYELOID LEUKEMIA

ONCOLOGY WATCH

New Year's Resolutions – A Different Perspective
Tempero M

COMMENTARY

Financial Toxicity in Patients With Cancer:
What a Tangled Web We Weave
Muluneh B

THE LAST WORD

Opioid and Benzodiazepine Use in Women
With Breast Cancer: More Work on the Horizon
Cata JP and Kim DD

MOLECULAR INSIGHTS IN PATIENT CARE

Targeting the *NTRK* Fusion Gene in
Pancreatic Acinar Cell Carcinoma:
A Case Report and Review of the Literature
Gupta M, Sherrow C, Krone ME, et al

REVIEWS

How Have Multigene Panels Changed the Clinical
Practice of Genetic Counseling and Testing
Pilarski R

Novel Agents for Metastatic Triple-Negative
Breast Cancer: Finding the Positive in the Negative
Vidula N, Ellisen LW, and Bardia A

ORIGINAL RESEARCH

New Persistent Opioid and Benzodiazepine
Use After Curative-Intent Treatment in
Patients With Breast Cancer
Sakamoto MR, Eguchi M, Azelby CM, et al

Impact of Axillary Dissection Among
Patients With Sentinel Node-Positive
Breast Cancer Undergoing Mastectomy
Sun J, Mathias BJ, Laronga C, et al

Gastrostomy Tube for Nutrition and Malignant
Bowel Obstruction in Patients With Cancer
Gauvin G, Do-Nguyen CC, Lou J, et al

Elderly Black Non-Hispanic Patients
With Head and Neck Squamous Cell Cancer
Have the Worst Survival Outcomes
McDermott JD, Eguchi M, Morgan R, et al

Patterns and Trends of Cancer Screening in Canada:
Results From a Contemporary National Survey
Abdel-Rahman O

New research in the January 2021 issue of *JNCCN—Journal of the National Comprehensive Cancer Network* finds more than a third of eligible people miss timely screening tests for colorectal cancer and at least a quarter appear to miss timely screening tests for breast and cervical cancers. The study comes from the University of Alberta, Faculty of Medicine and Dentistry in Alberta, Canada, with findings based on self-reported results from the Canadian Community Health Survey (CCHS) from 2007-2016. According to the author, the results also point to evidence of screening disparities being linked to lower socioeconomic status and identifiable minority race—echoing a similar study conducted in the United States by the Centers for Disease Control.

"We already have high-quality evidence showing that recommended [cancer screening](#) saves lives. From a cost perspective, it is far cheaper to have an efficient cancer screening program and lower cancer-specific mortality than to have a high incidence of advanced incurable cancers that would cost the system much more—not to mention the lost opportunities for healthy, cancer-free individuals to work and contribute to the economy," said Omar M. Abdel-Rahman Abdelsalam, MBBCh, MSc, MD, Assistant Professor of Medical Oncology, University of Alberta. "These findings show we need more measures to eliminate socioeconomic-related health disparities in our communities, even where healthcare is publicly funded. The most powerful intervention to improve screening rates would be to invest more in primary care and make sure every individual is linked to a family physician who can track their adherence to recommended screening tests."

Dr. Abdel-Rahman looked at self-reported data for 99,820 people eligible for [colorectal cancer](#) screening, 59,724 people eligible for breast

cancer screening, and 46,767 in the cervical cancer screening cohort, based on Canadian guidelines from 2007 through 2016. 43% did not have a timely screening test for colorectal, 35% did not have timely mammography, and 25% did not have a timely PAP smear. There were modest increases in screening compliance throughout the study's duration according to self-reporting within the survey.

"Dr. Abdel-Rahman's study adds to the literature showing that decreased levels of adherence to common cancer screening practices are associated with social and economic disparities," commented Mark Helvie, MD, Director, Breast Imaging Division, University of Michigan Rogel Cancer Center, Vice-Chair, NCCN Guidelines Panel for Breast Cancer Screening and Diagnosis, who was not involved with this research.

"Colorectal cancer screening had the lowest compliance with over a third reporting not being screened. While important differences exist between the NCCN Guidelines for screening and the Canadian screening guidelines—especially regarding colonoscopy and initiation and frequency of mammography—they all agree there is a need for more targeted efforts, research, and education to understand and improve compliance among all individuals, as illustrated by this study's results."

The research data does not include any impact from the COVID-19 pandemic. However, the publication is particularly timely as doctors report concerning drop-offs of screening numbers, which are projected to possibly result in thousands of excess cancer deaths over the next ten years. The study further highlights how economic and racial disparities contribute to unequal health outcomes and showcases the need for interventions that go beyond coverage for health services alone.

More information: Ingrid J. Hall et al, Patterns and Trends in Cancer Screening in the United States, *Preventing Chronic Disease* (2018). [DOI: 10.5888/pcd15.170465](https://doi.org/10.5888/pcd15.170465)

Provided by National Comprehensive Cancer Network

Citation: New research highlights dangerous disparities for life-saving cancer screening (2021, January 13) retrieved 6 May 2024 from <https://medicalxpress.com/news/2021-01-highlights-dangerous-disparities-life-saving-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.