


# Myeloid immune cells in the blood tied to severe COVID-19

January 26 2021

---



Anna Smed Sørensen, associate professor at the Department of Medicine, Solna, Karolinska Institutet. Credit: Ulf Sirborn

Individual variations in how the immune system responds to SARS-CoV-2 appear to impact the severity of disease. Researchers at

Karolinska Institutet in Sweden have now been able to show that patients with severe COVID-19 have significantly elevated levels of a certain type of immune cells in their blood, called myeloid-derived suppressor cells. The study published in the *Journal of Clinical Investigation* may bring an increased understanding of how early immune responses impact disease severity.

Most individuals with COVID-19 develop mild to moderate symptoms and recover without needing hospital treatment. In severe cases, however, COVID-19 can lead to respiratory failure or even death. It is not yet known why the severity of disease varies so much between patients.

Researchers at Karolinska Institutet, Karolinska University Hospital, Stemirna Therapeutics in Shanghai, and Stanford University in the United States have now studied one type of immune cell, monocytic myeloid-derived suppressor [cells](#), or M-MDSC, and their potential role in COVID-19.

## **Dampens T cell activity**

T cells are part of the [immune system](#) and play an important part in the body's protection against viral infections such as COVID-19. M-MDSCs have been shown to increase in other inflammatory conditions, and their suppressive effect on T cell activity has been established. The role of M-MDSC in respiratory infections, however, is largely unknown. Since low levels of T cells are a hallmark of COVID-19, it is of interest to understand the role of M-MDSCs in this disease.

The study consisted of 147 patients with mild to fatal COVID-19 who were sampled repeatedly from blood and the respiratory tract. These were then compared with patients with influenza and healthy individuals.

## Unbalanced immune system

The results show that patients with severe COVID-19 have significantly elevated levels of M-MDSCs in blood compared with milder cases and healthy individuals. COVID-19 patients had fewer T cells in blood than healthy subjects, and they showed signs of impaired function.

The analysis also showed that the levels of M-MDSCs early in the course of disease seemed to reflect subsequent disease severity.

"Our results help increase the understanding of what causes severe COVID-19 and is an important piece of the puzzle in understanding the connection between the early, innate immune system, which includes M-MDSC, and the later, adaptive immune system, which includes T cells. There is also a strong clinical connection, as you could potentially use the results to find new biomarkers for severe illness," says Anna Smed Sörensen, associate professor at the Department of Medicine, Solna, Karolinska Institutet, and the study's last author.

An inherent limitation of the study is the number of patients and amount of sample material that could be collected, why each sample was used as efficiently as possible.

"The next step in our research is to further study the connection between different parts of the immune system, such as M-MDSC, T cells, and antibodies," says Anna Smed Sörensen.

**More information:** Functional monocytic myeloid-derived suppressor cells increase in blood but not airways and predict COVID-19 severity.

[DOI: 10.1172/JCI44734](https://doi.org/10.1172/JCI44734).

Provided by Karolinska Institutet

Citation: Myeloid immune cells in the blood tied to severe COVID-19 (2021, January 26)  
retrieved 18 April 2024 from

<https://medicalxpress.com/news/2021-01-myeloid-immune-cells-blood-tied.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.