

More than half of people using cannabis for pain experience multiple withdrawal symptoms

January 8 2021



An illustration of withdrawal symptoms related to cannabis use. Credit: University of Michigan

More than half of people who use medical marijuana products to ease pain also experience clusters of multiple withdrawal symptoms when they're between uses, a new study finds.

And about 10% of the patients taking part in the study experienced worsening changes to their sleep, mood, [mental state](#), energy and appetite over the next two years as they continued to use [cannabis](#).

Many of them may not recognize that these symptoms come not from their underlying condition, but from their brain and body's reaction to the absence of substances in the cannabis products they're smoking, vaping, eating or applying to their skin, says the University of Michigan Addiction Center psychologist who led the study.

When someone experiences more than a few such symptoms, it's called cannabis [withdrawal](#) syndrome—and it can mean a higher risk of developing even more serious issues such as a cannabis use disorder.

In the new research published in the journal *Addiction*, a team from the U-M Medical School and the VA Ann Arbor Healthcare System reports findings from detailed surveys across two years of 527 Michigan residents. All were participating in the state's system to certify people with certain conditions for use of medical cannabis, and had non-cancer-related pain.

"Some people report experiencing significant benefits from medical cannabis, but our findings suggest a real need to increase awareness about the signs of withdrawal symptoms developing to decrease the potential downsides of cannabis use, especially among those who experience severe or worsening symptoms over time," says Lara Coughlin, Ph.D., the addiction psychologist who led the analysis.

Long-term study in medical cannabis use

The researchers asked the patients whether they had experienced any of 15 different symptoms—ranging from trouble sleeping and nausea to irritability and aggression—when they had gone a significant time

without using cannabis.

The researchers used an analytic method to empirically group the patients into those who had no symptoms or mild symptoms at the start of the study, those who had moderate symptoms (meaning they experienced multiple withdrawal symptoms) and those who had severe withdrawal issues that included most or all of the symptoms.

They then looked at how things changed over time, surveying the patients one year and two years after their first survey.

At baseline, 41% of the study participants fell into the mild symptoms group, 34% were in the moderate group and 25% were classed as severe.

Misconceptions about medical cannabis

Many people who turn to medical cannabis for pain do so because other pain relievers haven't worked, says Coughlin, an assistant professor in the Department of Psychiatry who sees patients as part of U-M Addiction Treatment Services. They may also want to avoid long-term use of opioid pain medications because they pose a risk of misuse and other adverse health consequences.

She notes that people who experience issues related to their cannabis use for pain should talk with their health care providers about receiving other pain treatments including psychosocial treatments such as cognitive behavioral therapy.

The perception of cannabis as "harmless" is not correct, she says. It contains substances called cannabinoids that act on the brain—and that over time can lead the brain to react when those substances are absent.

In addition to a general craving to use cannabis, withdrawal symptoms

can include anxiety, sleep difficulties, decreased appetite, restlessness, depressed mood, aggression, irritability, nausea, sweating, headache, stomach pain, strange dreams, increased anger and shakiness.

Previous research has shown that the more symptoms and greater severity of symptoms a person has, the less likely they are to be able to reduce their use of cannabis, quit using it or stay away from it once they quit.

They may mistakenly think that the symptoms happen because of their underlying medical conditions, and may even increase the amount or frequency of their cannabis use to try to counteract the effect—leading to a cycle of increasing use and increasing withdrawal.

Coughlin says people who decide to use a cannabis product for a medical purpose should discuss the amount, route of administration, frequency and type of cannabis product with their regular health provider. They should also familiarize themselves with the symptoms of cannabis withdrawal and tell their provider if they're experiencing them.

Feeling the urge to use cannabis after a period without use, such as soon after waking up, can be a sign of a withdrawal syndrome, she notes. So can the inability to cut back on use without experiencing craving or other symptoms of withdrawal.

Because there is no medically accepted standard for medical cannabis dosing for different conditions, patients are often faced with a wide array of cannabis products that vary in strength and route of administration. Some products could pose more risk for development of withdrawal symptoms than others, Coughlin says.

For example, people who smoked cannabis tended to have more severe withdrawal symptoms than others, while people who vaped cannabis

reported symptoms that tended to stay the same or get worse, but generally did not improve, over time.

As more states legalize cannabis for medical or general use, including several states that will legalize its use based on the results of last November's election, use is expected to grow.

More about the study

The researchers asked the patients about how they used [cannabis products](#), how often, and how long they'd been using them, as well as about their mental and physical health, their education and employment status.

Over time, those who had started off in the mild withdrawal [symptom](#) group were likely to stay there, but some did progress to moderate withdrawal symptoms.

People in the moderate withdrawal group were more likely to go down in symptoms than up, and by the end of the study the number of the people in the severe category had dropped to 17%. In all, 13% of the patients had gone up to the next level of symptoms by the end of the first year, and 8% had transitioned upward by the end of two years.

Sleep problems were the most common symptom across all three groups, and many in the mild group also reported cravings for cannabis. In the moderate group, the most common withdrawal symptoms were sleep problems, depressed mood, decreased appetite, craving, restlessness, anxiety and irritability.

The severe withdrawal symptom group was much more likely to report all the symptoms except sweatiness. Nearly all the participants in this group reported irritability, anxiety, and sleep problems. They were also

more likely to be longtime and frequent users of cannabis.

Those in the severe group were more likely to be younger and to have worse mental health. Older adults were less likely to go up in withdrawal symptom severity, while those who vaped cannabis were less likely to transition to a lower withdrawal-severity group.

The study didn't assess nicotine use, or try to distinguish between symptoms that could also be related to breakthrough pain or diagnosed/undiagnosed mental health conditions during abstinence.

Future directions

Coughlin and her colleagues hope future research can explore cannabis withdrawal symptoms among medical cannabis patients further, including the impact of different attempts to abstain, different types of use and administration routes, and interaction with other physical and mental health factors. Most research on cannabis withdrawal has been in recreational users, or "snapshot" looks at [medical cannabis](#) patients at a single point in time.

Further research could help identify those most at risk of developing problems, and reduce the risk of progression to cannabis use disorder, which is when someone uses cannabis repeatedly despite major impacts on their lives and ability to function.

More information: Lara N. Coughlin et al, Progression of cannabis withdrawal symptoms in people using medical cannabis for chronic pain, *Addiction* (2021). [DOI: 10.1111/add.15370](https://doi.org/10.1111/add.15370)

Provided by University of Michigan

Citation: More than half of people using cannabis for pain experience multiple withdrawal symptoms (2021, January 8) retrieved 26 April 2024 from <https://medicalxpress.com/news/2021-01-people-cannabis-pain-multiple-symptoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.