

Better post-surgery care would dramatically improve cancer survival

January 21 2021



Credit: Pixabay/CC0 Public Domain

Patients in low-and lower-middle-income countries were up to six times more likely to die from complications within 30 days of surgery compared with those in high-income countries, the report said.

Hospitals in these countries were found to be less likely to have post-surgical facilities or care plans in place.

Investing in appropriate recovery and ward space, trained staff, early warning systems and critical care facilities would result in much improved surgical care and reduce the number of deaths, experts said.

In the largest study of its kind, researchers from the Universities of Edinburgh and Birmingham examined data for nearly 16,000 patients in 428 hospitals across 82 countries who underwent surgery for breast, bowel and [stomach cancer](#) between April 2018 and January 2019.

Surgery is an important part of [cancer](#) treatment with 80 percent of cancer patients undergoing a procedure.

Stomach cancer patients who underwent surgery were three times more likely to die in low and lower-middle income countries than those in [high-income countries](#).

It was four times more likely for bowel [cancer patients](#) in low and lower-middle-income countries to die than those in high-income countries.

There was no difference in deaths between countries for patients who underwent breast cancer surgery.

Complications following surgery are common, but hospitals that provide a high standard of post-operative care had the best outcomes, even when treating late stage cancers, the study found.

Low- and lower-middle-income-countries that had post-operative care facilities in place were associated with seven to ten fewer deaths per 100 complications.

The research, published in *The Lancet*, has been funded through the National Institute for Health Research Global Health Unit in Global Surgery.

The team only looked at early outcomes following surgery, but, in future, they plan to study longer-term outcomes and other cancers.

Professor Ewen Harrison, Professor of Surgery and Data Science, University of Edinburgh, said: "Rich and poor countries alike have talented surgeons and anaesthesiologists, but low resource countries do not have the infrastructure to support the complications that occur during [surgery](#). We now know this can have a major impact on whether or not a patient survives."

More information: *The Lancet* (2021). [DOI: 10.1016/S0140-6736\(21\)00001-5](https://doi.org/10.1016/S0140-6736(21)00001-5)

Provided by University of Edinburgh

Citation: Better post-surgery care would dramatically improve cancer survival (2021, January 21) retrieved 26 June 2024 from <https://medicalxpress.com/news/2021-01-post-surgery-cancer-survival.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--