

# Severe menopause symptoms often accompany premature ovarian insufficiency

January 20 2021

---

Hot flashes, insomnia, and vaginal dryness are commonly reported symptoms that accompany the menopause transition. A new study suggests that such symptoms—especially psychological and sexual problems—are worse for women who have premature ovarian insufficiency (POI) than for women undergoing natural menopause. Study results are published online today in *Menopause*, the journal of The North American Menopause Society (NAMS).

Premature ovarian insufficiency is defined as the cessation of ovarian function that leads to menopause before the age of 40 years. The condition is associated with increased risks for a number of long-term health comorbidities, including osteoporosis, cardiovascular disease, [mood disorders](#), cognitive impairment, and sexual dysfunction. It can also lead to a shorter life expectancy.

Despite research around the various risk factors associated with POI, few studies have examined the effect of POI relative to the prevalence, severity, and factors affecting menopause symptoms. In this new study involving nearly 300 Chinese women, researchers specifically investigated menopause symptoms in women with POI and compared them with the severity and prevalence of similar symptoms in women who experienced natural menopause.

What they found was that women with POI experience a high prevalence of menopause symptoms, especially those in the psychological and sexual domains, and that these symptoms are often more severe than

those experienced by women who undergo natural menopause. The symptoms likely to be most severe include [mood swings](#), hot flashes, insomnia, fatigue, and sexual dysfunction (including [vaginal dryness](#), dyspareunia, and decreased libido).

On the basis of these results, the researchers have suggested that healthcare providers be more cognizant of psychological complications and sexual dysfunction risks in women with POI and work with them to identify options for relief.

Results are published in the article "Menopausal symptoms in women with premature ovarian insufficiency: prevalence, severity and associated factors."

"This study of Chinese women showed that those with POI had more prevalent menopause-related symptoms than women with menopause at the average age. It is important to note that regardless of the prevalence or severity of symptoms, women with POI should receive hormone therapy at least until the average age of menopause unless there are contraindications to its use in order to mitigate potential long-term adverse health effects, including fractures, heart disease, cognitive impairment, and early mortality—in addition to mood disturbances and [sexual dysfunction](#)," says Dr. Stephanie Faubion, NAMS medical director.

Provided by The North American Menopause Society

Citation: Severe menopause symptoms often accompany premature ovarian insufficiency (2021, January 20) retrieved 6 May 2024 from <https://medicalxpress.com/news/2021-01-severe-menopause-symptoms-accompany-premature.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.