


# Simple monitoring could reduce medicine misuse in care homes

January 11 2021

---



Credit: Unsplash/CC0 Public Domain

New research from Swansea University suggests that a simple nurse- or carer-led medicines' monitoring system can help reduce medication-related illness for people living in residential care homes—and the

process takes just a few minutes per patient.

The [research paper](#) published in the *PLOS ONE* journal looked how the monitoring system, known as the Adverse Drug Reaction Profile (ADRe-p), can help nurses or carers identify medicines' mismanagement or [adverse drug reactions](#) in patients prescribed multiple medicines, and can help avoid medication-related harm and improve prescribing.

Professor Sue Jordan, who led the study said: "The problem presented by the scale and complexity of inadvertent harm from both use and misuse of medicines is very real, which is reflected in the World Health Organisation's (WHO) Third Global Patient Safety Challenge aiming to reduce avoidable medication-related harm by 50% by next year.

"Our study took place in three independent private sector registered care homes. All the homes were very good, and we witnessed excellent nursing care. Nevertheless, ADRe helped to improve medicines management. Nurses or carers using the ADRe monitoring system identified possible medication-related harms and, following a review by doctors or pharmacists, new medication regimens were introduced.

The outcomes for the 19 patients involved meant that:

- 6 residents were no longer in pain
- 3 no longer experienced convulsions
- 3 no longer experienced aggression
- 2 had swallowing difficulties treated
- 4 no longer reported insomnia
- 1 had breathing difficulties treated
- 2 had their laxative prescriptions adjusted to reduce diarrhea
- falls ceased for 2 residents (of 4 noted as falling and of 5 able to stand).

The research team also found that few new problems arose, there was no clinical deterioration, and no harms were associated with the intervention.

Professor Jordan said: "What was really important about this study was that it showed that just 10 minutes of nurses' or carers' time, along with 10 minutes for a pharmacist review, made a huge difference to patient wellbeing. The use of ADRe not only improved residents' health, but also changed prescription regimens to ensure the maximum clinical benefits for patients, which ultimately helps to optimize healthcare resources."

**More information:** Sue Jordan et al, Nobody ever questions—Polypharmacy in care homes: A mixed methods evaluation of a multidisciplinary medicines optimisation initiative, *PLOS ONE* (2021). [DOI: 10.1371/journal.pone.0244519](https://doi.org/10.1371/journal.pone.0244519)

Provided by Swansea University

Citation: Simple monitoring could reduce medicine misuse in care homes (2021, January 11) retrieved 26 April 2024 from <https://medicalxpress.com/news/2021-01-simple-medicine-misuse-homes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--