

Early behavioural problems predict adolescent mental health difficulties

February 11 2021



Credit: Unsplash/CC0 Public Domain

A substantial proportion of adolescent mental health and behavioral difficulties can be predicted years before they arise, a new study indicates.

The research, by academics at the University of Cambridge and Royal Holloway, University of London, shows that children who experience certain [behavioral challenges](#), like hyperactivity and anxiety, are more likely to go on to develop poor mental health in adolescence, such as [emotional difficulties](#).

It also finds that children with behavioral difficulties are more likely to experience problems as adolescents if they come from less affluent, or disadvantaged backgrounds.

Levels of [adolescent](#) behavioral and mental health difficulties are rising. As healthcare systems worldwide struggle to cope with increasing demand, there is an urgent need to predict which adolescents are most likely to need additional support as early as possible.

The researchers used [historical data](#) from more than 6,700 people, collected both when they were 10 and 16 years old. For the first time, this allowed them to plot previously-undocumented connections between a wide range of childhood problems, such as hyperactivity and anxiety, and related but different behavioral difficulties that arise in adolescence.

They combined this information with details about the participants' social and economic backgrounds. Children from wealthier or better-resourced settings were more likely to overcome [behavioral problems](#) by the age of 16. Conversely, those from poorer backgrounds were more likely to transition into different difficulties during their teens.

Dr. Duncan Astle, from the MRC Cognition and Brain Sciences Unit, University of Cambridge, said: "At present, adolescent behavior and mental health difficulties are only treated once they become problematic, which is one of the main reasons why mental health services are overwhelmed. Our work shows how we might begin to develop a way to predict the difficulties of some—perhaps

many—young people and intervene sooner."

Professor Anna Vignoles, who co-authored the research while based at the Faculty of Education, University of Cambridge, said: "This type of information is essential if we want a more proactive, preventative model of handling behavioral and [mental health problems](#) in young people. The more significant predictors we have, the better we can target support."

The historical data came from the British Cohort Study—an ongoing project which is following the lives of the same group of people born during one week in April 1970. It contains details about behavioral difficulties the participants displayed both at 10 and 16 years old, as well as information about their socio-economic circumstances, such as [family income](#), and their parents' level of education. Few other sources offer such rich data on that scale, which is essential to making a robust assessment of how behavioral problems may change during a person's lifetime.

The researchers described each participant in the study at both 10 and 16 using six broad categories of potential behavioral problems: conduct problems, hyperactivity and impulsivity, inattention, emotional control problems, anxiety, and motor problems.

Next, they used a technique called hybrid hierarchical clustering which gathered together groups of similar profiles in a manner designed to offer the most meaningful interpretation of the data possible. In many cases, children fell into the 'no problems' group; some displayed one specific problem; while others presented combined patterns of difficulties (for example, one group of children exhibited a similar pattern of motor, hyperactivity and emotional problems combined).

The profile groups which emerged were very different across the two developmental stages. Hyperactivity, motor control and conduct issues

were prominent in childhood; while adolescents were more likely to display problems related to emotional control, anxiety and inattention.

Dr. Joe Bathelt, Lecturer at Royal Holloway, University of London, said: "We often assume that behavioral problems stay the same across development; however, our results show that the presentation can change substantially between childhood and adolescence."

The researchers then looked for cases where a larger-than-expected number of children had transitioned from one specific sub-group at age 10 to another aged 16.

Of those who experienced behavioral problems as children, 55% exhibited no difficulties in adolescence. There were, however, several cases of a clear relationship between specific clusters of childhood and adolescent problems. The main patterns that emerged were:

- An unexpectedly high percentage of children with anxiety problems (22%) developed emotional problems during adolescence.
- Around a quarter (24%) of children with conduct problems displayed 'a constellation of problems' with anxiety, emotion and inattention in adolescence.
- Around 17% of children with combined emotion, motor and hyperactivity problems developed inattention problems in adolescence.

When this data was combined with the participants' demographic context the researchers also found that the 55% of children with behavioral problems who had no problems by their mid-teens typically came from wealthier families, with more educated parents who tended to have higher-prestige jobs. They also tended to have higher cognitive ability, pointing to better educational attainment.

In addition, the data showed clearly that children who developed emotional control [problems](#) during adolescence were significantly more likely to come from poorer backgrounds, typically had more siblings, and had parents with lower-prestige jobs.

"It may be that wealthier families have more resources at their disposal and can seek help for [children](#) with behavioral disorders," Vignoles said. "Equally, growing up in an unstable environment is a significant predictor of the onset of adolescent behavioral difficulties, and that may be reflected here."

The researchers hope to carry out further studies which will compare the patterns that they recorded with data gathered from more recent generations. "It would be really interesting to rerun this analysis with the Millenium Cohort Study of people born in 2000," Astle said. "We need to bear in mind that what it means to be an adolescent has changed substantially over time, and will probably continue to do so."

More information: *Social Psychiatry and Psychiatric Epidemiology*,
[DOI: 10.1007/s00127-020-02014-4](https://doi.org/10.1007/s00127-020-02014-4)

Provided by University of Cambridge

Citation: Early behavioural problems predict adolescent mental health difficulties (2021, February 11) retrieved 4 May 2024 from <https://medicalxpress.com/news/2021-02-early-behavioural-problems-adolescent-mental.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
