

Fetal surgery for spina bifida leads to better mobility in school-age children

February 8 2021



Credit: Pixabay/CC0 Public Domain

Adding to a growing body of research affirming the benefits of fetal surgery for spina bifida, new findings show prenatal repair of the spinal column confers physical gains that extend into childhood. The



researchers found that children who had undergone fetal surgery for myelomeningocele, the most severe form of spina bifida, were more likely than those who received postnatal repair to walk independently, go up and down stairs, and perform self-care tasks like using a fork, washing hands and brushing teeth. They also had stronger leg muscles and walked faster than children who had their spina bifida surgery after birth.

The findings were published today in JAMA Pediatrics.

"This study shows that the benefits of fetal surgery for spina bifida extend beyond early childhood and well into a child's first decade of life," said N. Scott Adzick, MD, Surgeon-in-Chief at Children's Hospital of Philadelphia (CHOP), Director of CHOP's Center for Fetal Diagnosis and Treatment, and co-author of the study. "This is especially important because of previously raised concerns that the advantages from fetal surgery may decrease over time. Contrary to those concerns, there appears to be a long-term benefit from neural protection in utero."

Spina bifida is a <u>birth defect</u> affecting up to 1 in 1,500 births in the United States and occurs when an area of the spinal column does not form properly, leaving a section of the spinal cord and spinal nerves exposed through an opening in the back. The condition often results in weakness or paralysis below the location of the defect, leading to an inability to walk unassisted and a loss of bladder and bowel control for many children.

MOMS2

To assess the long-term physical outcomes of patients receiving fetal surgery for spina bifida, the researchers assessed participants from the landmark Management of Myelomeningocele Study (MOMS) study, which was co-led by investigators at CHOP, Vanderbilt University



Medical Center, and the University of California, San Francisco, along with the data coordinating center at the George Washington University Biostatistics Center and published in 2011 in the *New England Journal of Medicine*. That study compared outcomes of prenatal and traditional postnatal repair of myelomeningocele at 12 and 30 months and demonstrated that prenatal repair can offer significant benefits. Babies with spina bifida who received fetal surgery were less likely to need a shunt for the buildup of spinal fluid in the brain. They were also better able to walk two and a half years after surgery than those operated on after birth and had better overall motor function.

The researchers studied almost all of the children from MOMS when they were school-aged and in 2020 published the overall results of MOMS2: Follow up of the Management Of Myelomeningocele Study in Pediatrics. Focusing primarily on neurocognitive outcomes as well as motor skills, the researchers found that up to ten years after surgery, children who had received prenatal repair showed better gross and <u>fine</u> motor skills. They also continued to need fewer shunts, surgeries and catheterizations and reported better quality of life. In separate analyses they demonstrated better control over their bladder and bowel movements; children who underwent fetal surgery were almost six times as likely to go to the bathroom on their own than those operated on after birth.

Additional Benefits to Fetal Surgery

In MOMS2, the researchers evaluated the children at a comprehensive study visit when the child was 5-10 years old. Children underwent a physical exam and functioning assessment of motor, self-care, and recreational skills. Patients were assessed for physical skills like being able to sit upright independently and, for those who could walk, being able to perform skills like jumping, walking up and down stairs, and standing on one foot.



Of the 183 families enrolled in the MOMS trial, 154 children received the physical functioning assessment as part of MOMS2. Those in the fetal surgery group performed a greater percentage of age-appropriate self-care skills (90.8%) compared to children in the postnatal repair group (85.5%) and were better able to complete tasks like using a fork, brushing teeth, and washing and drying hands. Children who received fetal surgery were also 70% more likely to walk independently, at higher speeds and with better walking style, and were more adept at advanced motor skills like walking up and down stairs compared with children in the postnatal repair group. Those in the fetal surgery group were also more than twice as likely to walk without braces on their legs compared with children in the postnatal repair group.

"These data are important for demonstrating that fetal surgery for spina bifida improves mobility well into school age, but the implications of these results are even more profound," said first author Amy J. Houtrow, MD, Ph.D., MPH, Pediatric Rehabilitation Medicine Division Chief at UPMC Children's Hospital of Pittsburgh. "When children are able to move around independently and interact with their peers, it dramatically changes the way they engage with the world. Better mobility makes socializing easier and improves their overall quality of life. Additionally, demonstrating more independence with self-care skills at school-age means they are more likely to be independent as adults."

Continued advances in fetal surgery for spina bifida

As a pioneer in the field, CHOP began performing fetal surgery for <u>spina bifida</u> in 1998, three years after CHOP launched the Center for Fetal Diagnosis and Treatment. Since then, the Center has continued to improve the technique and publish on the practice, leading to significant advances.

"When we began performing fetal surgery more than two decades ago,



we did so with the hope that the procedure would improve lives for <u>children</u> and their families," Dr. Adzick said. "As we continue to improve the technique, shortening <u>surgery</u> times and increasing the gestational age at birth, we are heartened by these results, which show the lasting benefits of <u>fetal surgery</u>."

More information: Houtrow AJ, MacPherson C, Jackson-Coty J, Rivera M, Flynn L, Burrows PK, Adzick NS, Fletcher JM, Gupta N, Howell LJ, Brock III JW, Lee H, Walker WO, Thom EA. "Prenatal repair and physical functioning among children with myelomeningocele: A follow-up to a Randomized Controlled Trial," *JAMA Pediatrics*, February 8, 2020. DOI: 10.1001/jamapediatrics.2020.5674

Provided by Children's Hospital of Philadelphia

Citation: Fetal surgery for spina bifida leads to better mobility in school-age children (2021, February 8) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2021-02-fetal-surgery-spina-bifida-mobility.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.