

High smoking dependence linked to depression

February 25 2021



Credit: CC0 Public Domain

An international study coordinated by the University of Helsinki provides new information on the connection between symptoms of depression and smoking dependence. The results support the importance of understanding the reasons behind smoking dependence to be able to help depressed smokers to quit, and thus possibly relieve depression

symptoms.

It has been widely reported that [smoking](#) is more common among the people suffering from [depression](#) than in the population in general. The actual mechanism of this association, however, remains unclear.

So far, studies about the association between smoking dependence and depression are very rare. This study, led by researchers from the University of Helsinki, focused on the different self-reported motivation factors for smoking. The work was carried out in collaboration with the Universities of Wisconsin and Missouri U.S., where the questionnaire was initially developed.

The study was based on more than 1400 Finnish twins who were [smokers](#) and provided smoking and depression information by responding to questionnaires. The results were published on February 24th in the esteemed scientific journal *Addiction*.

The study found that smokers who were more dependent were also more likely to be depressed. Of the different motives for smoking examined, the motives related to high levels of craving and automatic smoking behavior, as well as smoking to regulate [emotional states](#) were found to be the most strongly related to depression.

"Our results suggest that depressed persons are not smoking because of taste or taste-reward related to smoking. Rather, depression seems to be related to primary dependence motives and the regulation of mood," says Senior Researcher Maarit Piirtola, the first author of the study from the Institute for Molecular Medicine Finland at the University of Helsinki.

The unique twin study setting enabled the study team to rule out familial influences, including shared genetic vulnerability, by examining the association between depression and smoking dependence within twin

pairs, especially within identical pairs.

According to Senior Researcher Tellervo Korhonen, who led the study, it is important to understand the individual underlying reasons that maintain smoking dependence, so that tailored and multidisciplinary smoking cessation support can be provided.

Even though this study gives strong new evidence about the relation between smoking dependence and depressive symptoms, the researchers highlight that this was a cross-sectional analysis. "Therefore, we could not show which happens first: smoking dependence or depression among daily smokers," says Professor and Research Director Jaakko Kaprio from the University of Helsinki.

More information: Maarit Piirtola et al. The associations of smoking dependence motives with depression among daily smokers, *Addiction* (2021). [DOI: 10.1111/add.15390](https://doi.org/10.1111/add.15390)

Provided by University of Helsinki

Citation: High smoking dependence linked to depression (2021, February 25) retrieved 20 March 2024 from <https://medicalxpress.com/news/2021-02-high-linked-depression.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--