


Lessons from the flu season

February 11 2021



3D print of influenza virus. The virus surface (yellow) is covered with proteins called hemagglutinin (blue) and neuraminidase (red) that enable the virus to enter and infect human cells. Credit: NIH

Australian researchers have come up with two key recommendations from studies of the annual influenza season—one highlighting the benefits of antivirals in reducing repeat hospitalization, and the other to

watch for underlying cardiovascular disease.

While the world focuses on the rising COVID-19 death toll, seasonal influenza continues to cause significant [mortality](#) and poses a significant economic burden every year.

The South Australian study, conducted at two major metropolitan training hospitals between January 2016 and March 2020, collected data from 1,828 [adult patients](#) (average age around 66) who were hospitalized with influenza A and B.

Researchers compared outcomes for patients who did and did not take the antiviral oseltamivir within 48 hours of admission to the hospital..

The study in *International Journal of Infectious Diseases* describes how early administration of Oseltamivir appears to reduce 30-day readmissions and also reduce the length of hospital stay, although it did not reduce 30-day mortality.

"Our study builds a strong case for the use of this treatment in suspected influenza patients as early as possible ," says lead researcher Dr. Yogesh Sharma.

In 2017 there were 29,000 influenza-related hospitalisations, and 745 deaths due to influenza in Australia. However, social distancing and other regulations with COVID-19 has probably helped to reduce the burden on the hospital system last winter.

Flinders researcher Dr. Yogesh Sharma, who is a senior Flinders Medical Centre consultant and senior lecturer in Flinders University, says another study of the data, recently published in another international journal *IJC Heart & Vasculture*, could direct future management of influenza patients in another way.

"Our study of influenza-related hospitalisations was also useful in analyzing whether patients presenting with this respiratory illness were at higher risk of dying within 30 days if they had underlying cardiovascular disease issues," Dr. Sharma says.

Comparing flu patients' high-sensitive-troponins (hsTnT) [the specific markers of myocardial damage] upon admission, the study pointed to there being a case for using acutely high or chronic hsTnT levels to predict mortality risk within 30 days of discharge.

Of the 1,828 patients, troponin results were available for 617 (47.7%) patients. Of these, 62 (10%) had acute myocardial injury and 232 (37.6%) had chronic hsTnT elevation.

"Both inpatient and 30-day mortality were significantly higher among patients with acute (P

"When compared to patients with negative hsTnT, acute but not chronic hsTnT elevation was significantly associated with 30-day mortality after adjustment for various co-variates (HR 8.30, 1.80-17.84, P value = 0.013)."

As one of the largest studies of its kind in the world, these results could be used to improve health outcomes in hospitals around the world, Dr. Sharma says.

While flu is often mild to moderate in its symptoms, it may cause severe illness leading to hospitalization and even death.

Worldwide, seasonal [influenza](#) may cause between 291,243—645,832 (4.0-8.8 per 100,000 individuals) respiratory-related deaths annually, with two-thirds of the total mortality in individuals over 65 years of age.

More information: Yogesh Sharma et al, Effectiveness of Oseltamivir in reducing 30-day readmissions and mortality among patients with severe seasonal influenza in Australian hospitalized patients, *International Journal of Infectious Diseases* (2021). [DOI: 10.1016/j.ijid.2021.01.011](https://doi.org/10.1016/j.ijid.2021.01.011)

Provided by Flinders University

Citation: Lessons from the flu season (2021, February 11) retrieved 23 May 2024 from <https://medicalxpress.com/news/2021-02-lessons-flu-season.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--