

No links between opioids or certain antibiotics in pregnancy and major birth defects

February 10 2021



Credit: CC0 Public Domain

Two studies published by *The BMJ* today find no links between prescription opioids or macrolide antibiotics taken during pregnancy and

risk of major birth defects.

Both opioids (painkillers) and macrolides (antibiotics) are often prescribed during [pregnancy](#). Some previous studies have reported associations between exposure to opioids and macrolides and certain birth defects, but results are inconsistent.

Two studies published today set out to address this uncertainty.

The first study involved data for more than 82,000 US women who received two or more prescriptions for any opioid during early (first trimester) pregnancy from 2000 to 2015.

After taking account of other potential risk factors, the researchers found no clinically meaningful increase in the risk of major birth defects overall, or in heart malformations, club foot or [neural tube defects](#).

They did find a small increased risk for cleft palate (four to five additional cases of [cleft palate](#) per 10,000 pregnancies exposed to opioids in the first trimester) and say clinicians should advise patients about this potential risk.

But overall, they say these findings suggest that [prescription opioids](#) used in early pregnancy are not associated with a substantial increase in risk for most of the malformation types considered.

In the second study, researchers in Denmark compared data for nearly 1.2 million pregnancies from 1997 to 2016 and found no association between exposure to macrolides in early pregnancy and risk of major birth defects.

Infants born with major birth defects occurred in 35 per 1000 pregnancies exposed to macrolides compared with 37 per 1000

pregnancies exposed to penicillin and 33 per 1000 unexposed pregnancies.

Results were unchanged after further analysis of individual macrolides and specific birth defects according to [organ systems](#), suggesting that the findings are robust.

Both studies are observational, so can't establish cause, and they relied on women taking the drugs as prescribed. What's more, the researchers can't rule out the possibility that they may have missed some birth defects.

However, these were large studies using reliable medical insurance and national registry data, respectively, and were able to account for a range of other risk factors that may have influenced the results.

As such, both research teams say their findings provide reassurance about the risk of major [birth defects](#) when treatment with opioids or [macrolide antibiotics](#) are needed during pregnancy.

They also help inform clinicians, patients, and drug regulatory authorities about the selection of medications for women who are pregnant and women of reproductive age who may inadvertently become pregnant.

More information: Association of first trimester prescription opioid use with congenital malformations in the offspring: population based cohort study, *BMJ* (2021). www.bmj.com/content/372/bmj.n102

Association between use of macrolides in pregnancy and risk of major birth defects: nationwide, register based cohort study, *BMJ* (2021). www.bmj.com/content/372/bmj.n107

Provided by British Medical Journal

Citation: No links between opioids or certain antibiotics in pregnancy and major birth defects (2021, February 10) retrieved 6 May 2024 from <https://medicalxpress.com/news/2021-02-links-opioids-antibiotics-pregnancy-major.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.