


# Can I take painkillers before or after a COVID-19 vaccine?

February 4 2021, by Marilyn Marchione

---



Can I take painkillers before or after a COVID-19 vaccine? AP Illustration/Peter Hamlin

Can I take painkillers before or after a COVID-19 vaccine?

It's best to avoid them, unless you routinely take them for a [medical condition](#). Although the evidence is limited, some painkillers might interfere with the very thing the vaccine is trying to do: generate a strong

immune system response.

Vaccines work by tricking the body into thinking it has a virus and mounting a defense against it. That may cause arm soreness, fever, headache, muscle aches or other temporary symptoms of inflammation that can be part of that reaction.

"These symptoms mean your [immune system](#) is revving up and the vaccine is working," Dr. Rochelle Walensky, director of the U.S. Centers for Disease Control and Prevention, said in a recent news briefing.

Certain painkillers that target inflammation, including ibuprofen (Advil, Motrin and other brands) might curb the immune response. [A study on mice](#) in the *Journal of Virology* found these drugs might lower production of antibodies—helpful substances that block the virus from infecting cells.

If you're already taking one of those medications for a health condition, you should not stop before you get the vaccine—at least not without asking your doctor, said Jonathan Watanabe, a pharmacist at the University of California, Irvine.

People should not take a painkiller as a [preventive measure](#) before getting a [vaccine](#) unless a doctor has told them to, he said. The same goes for after a shot: "If you don't need to take it, you shouldn't," Watanabe said.

If you do need one, acetaminophen (Tylenol) "is safer because it doesn't alter your [immune response](#)," he added.

The CDC [offers other tips](#), such as holding a cool, wet washcloth over the area of the shot and exercising that arm. For fever, drink lots of

fluids and dress lightly.

Call your doctor if redness or tenderness in the arm increases after a day or if side effects don't go away after a few days, the CDC says.

**More information:** [Can the coronavirus travel more than 6 feet in the air?](#)

[How long could I be contagious before a positive virus test?](#)

© 2021 The Associated Press. All rights reserved. This material may not be published, broadcast, rewritten or redistributed without permission.

Citation: Can I take painkillers before or after a COVID-19 vaccine? (2021, February 4)  
retrieved 19 April 2024 from  
<https://medicalxpress.com/news/2021-02-painkillers-covid-vaccine.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--