

Peanut allergy affects even more U.S. adults than children

February 9 2021



Credit: CC0 Public Domain

Peanut allergy affects at least 4.5 million adults in the U.S., many of whom report developing their first allergy symptoms during adulthood, reports a new Northwestern Medicine study.

However, despite the fact that roughly three out of four Americans with [peanut allergy](#) are over 17 years old, [peanut allergy](#) is often considered a predominantly pediatric concern. For example, earlier this year the U.S. Food and Drug Administration (FDA) approved a peanut allergy therapy for initiation in pediatric patients ages 4-17. There are currently no FDA-approved therapies for patients with adult-onset food allergy.

The new study provides the first detailed estimates of peanut allergy among U.S. adults in all 50 states, which was previously unknown. These data indicate peanut allergy may be more common than previously acknowledged, and while younger adults are most affected, peanut allergy impacts U.S. adults of all ages.

"Currently, the sole FDA-approved peanut allergy therapy—Palforzia—is only indicated for pediatric patients. Given the high prevalence of peanut allergy among U.S. adults, additional therapies are needed to help address this growing burden of disease," said senior study author Dr. Ruchi Gupta, professor of pediatrics at Northwestern University Feinberg School of Medicine and a physician at Ann & Robert H. Lurie Children's Hospital of Chicago.

Gupta also is director of the Center for Food Allergy and Asthma Research at Feinberg.

The paper will be published Feb. 9 in the *Journal of Allergy and Clinical Immunology*.

The study found 2.9% of U.S. adults report a current peanut allergy, while 1.8% report a physician-diagnosed peanut allergy and/or a history of peanut-allergic reaction symptoms. In addition, researchers found two in three adults with peanut allergy have at least one other food allergy—most commonly tree nuts but more than one in five are also allergic to shellfish.

The data also indicate many individuals who report peanut allergies and experience potentially severe allergic reactions are not obtaining clinical diagnosis of their allergies.

"Clinical confirmation of suspected food allergies, no matter when their reported onset, is critical to reduce the risk of unnecessary allergen exposure as well as to ensure patients receive essential counseling and prescription of emergency epinephrine," said study co-first author Dr. Dawn Lei, a clinical instructor of pediatrics at Feinberg and an allergist and immunologist at Edward Hines Jr. VA Hospital.

Christopher Warren, director of population health at Feinberg's Center for Food Allergy and Asthma Research and co-first author added, "Unlike allergies such as milk or egg, which often develop early in life and are outgrown by adolescence, peanut allergy appears to affect children and adults to a similar degree. Our study shows many adults are not outgrowing their childhood peanut allergies, and many adults are developing peanut allergies for the first time.

"Worryingly, despite reporting similar rates of severe reactions and annual food allergy-related emergency room visits, patients with adult-onset peanut allergy were less likely to report a physician-diagnosis as well as a current epinephrine prescription."

Together, these data suggest additional efforts are needed to ensure optimal [food](#) allergy diagnosis and management among [adults](#) with peanut allergy, study authors said.

Provided by Northwestern University

Citation: Peanut allergy affects even more U.S. adults than children (2021, February 9) retrieved 23 April 2024 from

<https://medicalxpress.com/news/2021-02-peanut-allergy-affects-adults-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.