

How we sleep and experience psychological symptoms during pandemic

February 11 2021



Credit: Unsplash/CC0 Public Domain

A study shows that during the first confinement, day-to-day variations in subjective sleep quality influenced the occurrence of mental and physical health complaints, and that these effects were linked to daily reports of COVID-19 related deaths. The team of researchers led by

Peter Simor, CRCN -Center for Research in Cognition and Neurosciences and UNI, ULB Neurosciences Institute, Université libre de Bruxelles (ULB), interviewed 166 participants in three European countries, twice a day for two consecutive weeks via an online interface about their sleep quality and negative psychological experiences.

During the first confinement (18 March to 10 May 2020), people who reported worse [sleep quality](#) during a night also reported an increase in negative mood, psychotic-like experiences and somatic complaints on the next day. Furthermore, daily reports of deaths caused by COVID-19 predicted psychological symptoms on the same day and sleep quality the following night. This is the result of research carried out in three countries (Belgium, Hungary, Spain) under direction of Peter Simor with researchers at the Université libre de Bruxelles (ULB), Rebeca Sifuentes-Ortega, Ariadna Albajara Saenz, Oumaïma Benkirane, Anke Van Roy and Philippe Peigneux from the CRCN (Center for Research in Cognition and Neurosciences) and the UNI (ULB Neurosciences Institute) in collaboration with researchers from the Eötvös Loránd University of Budapest and the Universidad Autónoma de Madrid.

The quality of sleep, according to the researchers, could be related to the COVID-19 pandemic and even influence [mental health complaints](#). To test this hypothesis researchers interviewed 166 participants twice a day (in the morning and in the afternoon) for two consecutive weeks via an online interface about their sleep quality and negative psychological experiences. This method made it possible to measure variables (subjective sleep quality and daytime experiences such as rumination, psychotic-like experiences and somatic complaints) in a "prospective" manner and to establish general trends in individuals.

They were able to study the temporal associations of these variables within individuals, examining whether:

- a relatively poor night's sleep would be followed by more mental [health](#) complaints the next day.
- whether a relative increase in mental health complaints on a given day would lead to an altered sleep pattern the following night.

The research team also linked the measures (sleep and mental health complaints) to specific factors related to the COVID-19 pandemic: Daily COVID19 deaths reported by the national media were used as indicators of the threatening context of the pandemic. Thereby, they were able to show that daily media reports of the number of deaths related to COVID19 per country predicted mental health complaints that day and impaired sleep quality the following night.

The results show that changes in sleep quality from one night to the next predict how individuals cope with the daily challenges of confinement the next day.

More information: Péter Simor et al. Home confinement during the COVID-19: day-to-day associations of sleep quality with rumination, psychotic-like experiences, and somatic symptoms, *Sleep* (2021). [DOI: 10.1093/sleep/zsab029](https://doi.org/10.1093/sleep/zsab029)

Provided by Université libre de Bruxelles

Citation: How we sleep and experience psychological symptoms during pandemic (2021, February 11) retrieved 24 April 2024 from <https://medicalxpress.com/news/2021-02-psychological-symptoms-pandemic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.