

# How women make choices about surgery to prevent ovarian cancer

February 11 2021

---

A study by researchers at Queen Mary University in London has investigated how women who are at high risk of ovarian cancer make choices about possible preventive surgery.

Women who have BRCA1 and BRCA2 [genetic mutations](#) have a 17-44% lifetime risk of ovarian [cancer](#). Scientists now believe that most ovarian cancers begin in the fallopian tubes, and so [women](#) at high risk can choose to undergo surgery to remove both their [fallopian tubes](#) and ovaries to reduce their risk. For those who have not yet reached menopause, there are [difficult decisions](#) around the order and timing of surgeries, as removing the ovaries will induce menopause.

The women interviewed were 24 participants in the UK PROTECTOR trial, in which removal of tubes as an initial step, and delayed removal of ovaries in a second surgery, is being offered as an alternative to standard surgery, which removes both at once.

Standard surgery, removing both the tubes and ovaries, induces menopause in women who have not yet reached this stage of life. The proposed alternative two step protocol delays the induced menopause caused by the removal of the ovaries, and therefore avoids menopause side effects.

The study found that women who were more concerned about cancer risk than about menopause preferred the standard surgery, but those who were more concerned about the detrimental impact of menopause chose

the two-step procedure.

Women managed in specialist familial cancer clinic settings felt that they received better quality care than those in non-specialist clinics. The authors hope that this study will help [health professionals](#) to improve [decision support](#) provided to women who carry these genetic mutations.

Author Professor Ranjit Manchanda from Queen Mary University of London said: "This study shows that a number of women prefer a two step operation to manage their ovarian cancer risk. It is important for women to be given different options so that they can choose what is best for them. Women are happier with their care in specialist family cancer clinics. We need to set up a system for women at higher risk of gynaecological cancers to access such centres, which are able to provide a more streamlined, holistic approach." ?

**More information:** Surgical decision making in premenopausal BRCA carriers considering risk reducing early-salpingectomy or salpingo-oophorectomy: a Qualitative Study. Gaba F, Goyal S, Marks D, Chandrasekaran D, Evans, O, Robbani S, Tyson C, Legood R, Saridogan E, McCluggage GW, Hanson H, Singh N, Evans DG, Menon U, Manchanda R, on behalf of the PROTECTOR team. *J Med Genet* 2020.

Provided by Queen Mary, University of London

Citation: How women make choices about surgery to prevent ovarian cancer (2021, February 11) retrieved 10 May 2024 from <https://medicalxpress.com/news/2021-02-women-choices-surgery-ovarian-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.