

New discoveries of deep brain stimulation put it on par with therapeutics

March 25 2021, by Laurie Fickman



In deep brain stimulation, electrodes are surgically implanted in the deep brain and electrical pulses are delivered at certain rates to control tremors and other disabling motor signs associated with Parkinson's disease. Credit: University of Houston



Despite having remarkable utility in treating movement disorders such as Parkinson's disease, deep brain stimulation (DBS) has confounded researchers, with a general lack of understanding of why it works at some frequencies and does not at others. Now a University of Houston biomedical engineer is presenting evidence in *Nature Communications Biology* that electrical stimulation of the brain at higher frequencies (>100Hz) induces resonating waveforms which can successfully recalibrate dysfunctional circuits causing movement symptoms.

"We investigated the modulations in local field potentials induced by electrical stimulation of the subthalamic nucleus (STN) at therapeutic and non-therapeutic frequencies in Parkinson's disease patients undergoing DBS surgery. We find that therapeutic high-frequency stimulation (130-180 Hz) induces high-frequency oscillations (~300 Hz, HFO) similar to those observed with pharmacological treatment," reports Nuri Ince, associate professor of biomedical engineering.

For the past couple of decades, <u>deep brain stimulation</u> (DBS) has been the most important therapeutic advancement in the treatment of Parkinson's disease, a progressive nervous system disorder that affects movement in 10 million people worldwide. In DBS, electrodes are surgically implanted in the deep brain and electrical pulses are delivered at certain rates to control tremors and other disabling motor signs associated with the <u>disease</u>.

Until now, the process to find the correct frequency has been time consuming, with it taking sometimes months to implant devices and test their abilities in patients, in a largely back and forth process. Ince's method may speed the time to almost immediate for the programming of devices at correct frequencies.

"For the first time, we stimulated the brain and while doing that we recorded the response of the <u>brain</u> waves at the same time, and this has



been a limitation over the past years. When you stimulate with electrical pulses, they generate large amplitude artifacts, masking the neural response. With our signal processing methods, we were able to get rid of the noise and clean it up," said Ince. "If you know why certain frequencies are working, then you can adjust the <u>stimulation</u> frequencies on a subject-specific basis, making therapy more personalized."

More information: Musa Ozturk et al. Electroceutically induced subthalamic high-frequency oscillations and evoked compound activity may explain the mechanism of therapeutic stimulation in Parkinson's disease, *Communications Biology* (2021). DOI: 10.1038/s42003-021-01915-7

Provided by University of Houston

Citation: New discoveries of deep brain stimulation put it on par with therapeutics (2021, March 25) retrieved 27 April 2024 from <u>https://medicalxpress.com/news/2021-03-discoveries-deep-brain-simulation-par.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.