


# More evidence of cognitive dysfunction in 'long haul' non-hospitalized COVID-19 patients

March 25 2021, by Bob Yirka

---



SARS-CoV-2 (shown here in an electron microscopy image). Credit: National Institute of Allergy and Infectious Diseases, NIH

A team of researchers at Northwestern University's Feinberg School of

Medicine has found more evidence of cognitive dysfunction in many "long haul," non-hospitalized COVID-19 patients. In their paper published in the journal *Annals of Clinical and Translational Neurology*, the group describes their study of medical data for non-hospitalized COVID-19 patients across 21 states who were experiencing long-term symptoms from the disease and what they learned from it.

As the pandemic has worn on, [media outlets](#) continue to publish statistics regarding the number of infections and number of deaths from COVID-19—less attention has been paid to those people who were harmed by their [infection](#) but lived to tell medical personnel about their experiences. In this new effort, the researchers sought to learn more about some of the long-term effects experienced by people who have been infected by the SARS-CoV-2 virus.

To learn more about the "long haulers," as they have come to be known, the researchers obtained data from [medical facilities](#) in 21 states for 100 patients who had COVID-19 but who had not been admitted to the hospital. Some were seen via telehealth; others in person, many at Northwestern Memorial Hospital in Chicago. The researchers noted approximately half of the data involved patients who had not been diagnosed with COVID-19 but who were believed to have been infected.

The researchers found that approximately 85% of those patients they studied were reported to have had at least four [neurological symptoms](#)—the most common were brain fog, headaches and numbness or tingling. Other [common symptoms](#) included loss of taste and/or smell, blurred vision, dizziness and ringing in the ears. They noted that many of the patients described such symptoms as coming and going, and most reported their symptoms lasting for several months—some for as long as nine.

The researchers also found that many of those patients they studied had

reported feeling depressed or anxious prior to their infection, which could suggest a heightened risk of long-haul symptoms for people with similar conditions. They also acknowledge that their study was small, but suggest it was large enough to show that long-haul patients may need more care than has been realized.

**More information:** Edith L Graham et al. Persistent neurologic symptoms and cognitive dysfunction in non-hospitalized Covid-19 "long haulers", *Annals of Clinical and Translational Neurology* (2021). [DOI: 10.1002/acn3.51350](https://doi.org/10.1002/acn3.51350)

© 2021 Science X Network

Citation: More evidence of cognitive dysfunction in 'long haul' non-hospitalized COVID-19 patients (2021, March 25) retrieved 6 May 2024 from <https://medicalxpress.com/news/2021-03-evidence-cognitive-dysfunction-haul-non-hospitalized.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--