


Evidence supports COVID hearing loss link, say scientists

March 22 2021



Credit: Unsplash/CC0 Public Domain

Hearing loss and other auditory problems are strongly associated with COVID-19 according to a systematic review of research evidence led by University of Manchester and NIHR Manchester Biomedical Research

Centre (BRC) scientists.

Professor Kevin Munro and Ph.D. researcher Ibrahim Almufarrij found 56 studies that identified an association between COVID-19 and auditory and vestibular problems.

They pooled data from 24 of the studies to estimate that the prevalence of [hearing loss](#) was 7.6%, tinnitus was 14.8% and vertigo was 7.2%.

They publish their findings in the *International Journal of Audiology*.

However, the team—who followed up their review carried out a year ago—described the quality of the studies as fair.

Their data primarily used self-reported questionnaires or [medical records](#) to obtain COVID-19-related symptoms, rather than the more scientifically reliable [hearing](#) tests.

Kevin Munro, Professor of Audiology at The University of Manchester and Manchester BRC Hearing Health Lead said: "There is an urgent need for a carefully conducted clinical and diagnostic study to understand the long-term effects of COVID-19 on the auditory system.

"It is also well-known that viruses such as measles, mumps and meningitis can cause hearing loss; little is understood about the auditory effects of the SARS-CoV-2 virus."

"Though this review provides further evidence for an association, the studies we looked at were of varying quality so more work needs to be done."

Professor Munro, is currently leading a year-long UK study to investigate the possible long-term impact of COVID-19 on hearing

among people who have been previously treated in hospital for the virus.

His team hope to accurately estimate the number and severity of COVID-19 related hearing disorders in the UK, and discover what parts of the auditory system might be affected

They will also explore the association between these and other factors such as lifestyle, the presence of one or more additional conditions and critical care interventions.

A recent study led by Professor Munro, suggested that more than 13 percent of patients who were discharged from a hospital reported a change in their hearing.

Ibrahim Almufarrij said: "Though the evidence is of varying quality, more and more studies are being carried out so the evidence base is growing. What we really need are studies that compare COVID-19 cases with controls, such as patients admitted to hospital with other health conditions.

"Though caution needs to be taken, we hope this study will add to the weight of scientific evidence that there is a strong association between COVID-19 and hearing problems."

Professor Munro added: "Over the last few months I have received numerous emails from people who reported a change in their hearing, or tinnitus after having COVID-19.

"While this is alarming, caution is required as it is unclear if changes to hearing are directly attributed to COVID-19 or to other factors, such as treatments to deliver urgent care."

More information: *International Journal of Audiology* ,

www.tandfonline.com/doi/full/10.1080/14467887.2021.1896793

Provided by Taylor & Francis

Citation: Evidence supports COVID hearing loss link, say scientists (2021, March 22) retrieved 18 April 2024 from

<https://medicalxpress.com/news/2021-03-evidence-covid-loss-link-scientists.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.