


# Factors that may predict next pandemic

March 30 2021, by Loren Smith

---



Zoonotic pathogen diversity. Red = high diversity, green = low diversity. Credit: Singh et al.

New modeling identifies human and human-influenced environmental factors that are associated with disease outbreaks, including in Australia.

Humans are creating or exacerbating the environmental conditions that could lead to further pandemics, new University of Sydney research finds.

Modeling from the Sydney School of Veterinary Science suggests pressure on ecosystems, climate change and [economic development](#) are key factors associated with the diversification of pathogens (disease-causing agents, like viruses and bacteria). This has potential to lead to [disease outbreaks](#).

The research, by Dr. Balbir B Singh, Professor Michael Ward, and Associate Professor Navneet Dhand, is published in the international journal, Transboundary and Emerging Diseases.

They found a greater diversity of zoonotic diseases (diseases transmitted between animals and humans) in higher income countries with larger land areas, more dense human populations, and greater forest coverage.

The study also confirms increasing population growth and density are major drivers in the emergence of zoonotic diseases. The global human population has increased from about 1.6 billion in 1900 to about 7.8 billion today, putting pressure on ecosystems.

Associate Professor Dhand said: "As the human population increases, so does the demand for housing. To meet this demand, humans are encroaching on wild habitats. This increases interactions between wildlife, domestic animals and human beings which increases the potential for bugs to jump from animals to humans."

"To date, such disease models have been limited, and we continue to be frustrated in understanding why diseases continue to emerge," said Professor Ward, an infectious diseases expert. "This information can help inform disease mitigation and may prevent the next COVID-19."

Other zoonotic diseases that have recently devastated human populations include SARS, avian (H5N1) and swine (H1N1) flu, Ebola and Nipah—a bat-borne virus.

## Factors predicting disease

The researchers discovered country-level factors predicting three categories of disease: zoonotic, emerging (newly discovered diseases, or those diseases that have increased in occurrence or occurred in new locations), and human.

- Zoonotic diseases: land area, human population density, and area of forest. Areas with high zoonotic disease diversity include Europe, North America, Latin America, Australia, and China.
- Emerging diseases: land area, human population density and the human development index. Areas with high emerging disease diversity include Europe, North America, Latin America, and India.
- Human diseases: high per capita health expenditure, mean annual temperature, land area, human [population](#) density, human development index and rainfall. Areas with high human disease diversity include North America, Latin America, China and India.

"Countries within a longitude of -50 to -100 like Brazil, developed countries like United States and dense countries such as India were predicted to have a greater diversity of emerging diseases," Professor Ward said.

The researchers also noted weather variables, such as temperature and rainfall, could influence the diversity of human diseases. At warmer temperatures, there tend to be more emerging pathogens.

Their analyses demonstrate that weather variables (temperature and rainfall) have the potential to influence pathogen diversity.

These factors combined confirm human development—including human-

influenced climate change—not only damages our environment but is responsible for the emergence of infectious diseases, such as COVID-19.

## Using data to prevent outbreaks

"Our analysis suggests sustainable development is not only critical to maintaining ecosystems and slowing [climate change](#); it can inform [disease](#) control, mitigation, or prevention," Professor Ward said.

"Due to our use of national-level data, all countries could use these models to inform their public health policies and planning for future potential pandemics."

**Methodology:** The authors used 13,892 unique pathogen–country combinations and 49 socioeconomic and environmental variables to develop this model. Information from 190 countries was analyzed using statistical models to identify drivers for emerging and zoonotic diseases.

**More information:** Balbir B. Singh et al. Geodemography, environment and societal characteristics drive the global diversity of emerging, zoonotic and human pathogens, *Transboundary and Emerging Diseases* (2021). [DOI: 10.1111/tbed.14072](https://doi.org/10.1111/tbed.14072)

Provided by University of Sydney

Citation: Factors that may predict next pandemic (2021, March 30) retrieved 4 May 2024 from <https://medicalxpress.com/news/2021-03-factors-pandemic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.