

Foodborne fungus impairs intestinal wound healing in Crohn's disease

March 11 2021



High magnification micrograph of Crohn's disease. Biopsy of esophagus. H&E stain. Credit: Nephron/Wikipedia

Eating is a dangerous business. Naturally occurring toxins in food and potentially harmful foodborne microbes can do a number on our intestines, leading to repeated minor injuries. In healthy people, such damage typically heals in a day or two. But in people with Crohn's



disease, the wounds fester, causing abdominal pain, bleeding, diarrhea and other unpleasant symptoms.

Researchers at Washington University School of Medicine in St. Louis and the Cleveland Clinic have discovered that a fungus found in foods such as cheese and processed meats can infect sites of intestinal damage in mice and people with Crohn's and prevent <u>healing</u>. Moreover, treating infected mice with <u>antifungal medication</u> eliminates the fungus and allows the <u>wounds</u> to heal.

The findings, published March 12 in the journal *Science*, suggest that antifungal drugs and dietary changes are potential new approaches to improving intestinal wound healing and reducing symptoms of Crohn's disease.

"We're not suggesting that people stop eating cheese and processed meat; that would be going far beyond what we know right now," said first author Umang Jain, Ph.D., an instructor in pathology & immunology at the School of Medicine. "What we know is that this foodborne fungus gets into inflamed, injured tissue and causes harm. We're planning to perform a larger study in people to figure out if there's a correlation between diet and the abundance of this fungus in the intestine. If so, it is possible dietary modulation could lower levels of the fungus and thereby reduce symptoms of Crohn's disease."





Cleveland Clinic-led research team found that levels of the yeast D. hansenii are higher in Crohn's disease patients, particularly abundant within chronically inflamed regions of the colon and small intestine, indicative of unhealed intestinal wounds. Credit: Cleveland Clinic

Crohn's is a subtype of inflammatory bowel disease. As the name suggests, it is driven by <u>chronic inflammation</u> in the <u>digestive tract</u> and primarily treated with immunosuppressive medications. Crohn's patients endure repeated cycles of gastrointestinal symptom flare-up and remission. During a flare, their digestive tracts are dotted with inflamed, open sores that can persist for weeks or even months.

To understand why intestinal ulcers take so long to heal in some people, Jain and senior author Thaddeus Stappenbeck, MD, Ph.D., formerly at Washington University and now at the Cleveland Clinic, studied mice whose intestines had been injured. By sequencing microbial DNA at the site of injury, they discovered that the fungus Debaryomyces hansenii was abundant in wounds but not in uninjured parts of the intestine.



People acquire the fungus through their food and drink, Jain said. D. hansenii is commonly found in all kinds of cheeses, as well as sausage, beer, wine and other fermented foods.



A Cleveland Clinic-led research team found that levels of the yeast D. hansenii are higher in Crohn's disease patients, particularly abundant within chronically inflamed regions of the colon and small intestine, indicative of unhealed intestinal wounds. Credit: Cleveland Clinic

Further experiments showed that introducing the fungus into mice with injured intestines slowed down the <u>healing process</u>, and that eliminating D. hansenii with the antifungal drug amphotericin B sped it up.

People with Crohn's disease carry the same fungus as the mice. Jain and Stappenbeck examined intestinal biopsies from seven people with Crohn's disease and 10 healthy people. All seven of the patients harbored the fungus in their gut tissue, as compared with only one of the healthy people. In a separate analysis of 10 Crohn's patients involving <u>tissue</u>



<u>samples</u> of both inflamed and uninflamed areas of the gut, the researchers found the fungus in samples from all of the patients but only at sites of injury and inflammation.

"If you look at stool samples from healthy people, this fungus is highly abundant," Jain said. "It goes into your body and comes out again. But people with Crohn's disease have a defect in the intestinal barrier that enables the fungus to get into the tissue and survive there. And then it makes itself at home in ulcers and sites of inflammation and prevents those areas from healing."

The findings suggest that eliminating the fungus might restore normal wound healing and shorten flare-ups. While the drug amphotericin B was effective at eliminating the fungus in mouse studies, it is not widely used in people because it can only be given intravenously. The researchers are working with chemists to develop an effective antifungal that can be taken by mouth. They also are studying whether there is a link between diet and the amount of the fungus in people's digestive tracts.

"Crohn's disease is fundamentally an inflammatory disease, so even if we figured out how to improve wound healing, we wouldn't be curing the disease," Jain said. "But in people with Crohn's, impaired wound healing causes a lot of suffering. If we can show that depleting this <u>fungus</u> in people's bodies—either by dietary changes or with antifungal medications—could improve wound healing, then it may affect the quality of life in ways that we've not been able to do with more traditional approaches."

More information: U. Jain el al., "Debaryomyces is enriched in Crohn's disease intestinal tissue and impairs healing in mice," *Science* (2021). <u>science.sciencemag.org/cgi/doi ... 1126/science.abd0919</u>



Provided by Washington University School of Medicine

Citation: Foodborne fungus impairs intestinal wound healing in Crohn's disease (2021, March 11) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2021-03-foodborne-fungus-impairs-intestinal-wound.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.