

Hip fracture outcomes worse during busy periods

March 10 2021, by Av Steinar Brandslet



Credit: Pixabay/CC0 Public Domain

Hip fractures are serious, especially for the elderly. The operation can be a great strain, and 13 percent of patients over the age of 70 do not survive 60 days after the fracture.

Their chance of survival may depend on how busy the surgeons are with other emergency procedures.

"When the [operating room](#) is busy, 20 percent more of the patients die within 60 days after the operation," says Professor Johan Håkon Bjørngaard at the Norwegian University of Science and Technology's (NTNU) Department of Public Health and Nursing.

Surgeons can get especially busy during periods when the patient demand for [surgery](#) is high. In busy periods, hip fracture patients have to wait on average 20 percent longer before being operated on compared to the quiet periods. This wait can have serious consequences.

Information from more than 60,000 hip surgeries and all simultaneous emergency surgeries provided the research group from St. Olavs Hospital and NTNU with a solid numerical basis.

"We investigated how many older people over the age of 70 died during the first 60 days following a hip operation when particularly many emergency patients were queued up for surgery at the hospitals," say researchers Andreas Asheim and Sara Marie Nilsen from the Regional Center for Health Services Development (RSHU) at St. Olavs [hospital](#).

During busy periods, 40 percent of the patients waiting in the operating wards are typically people who have recently been brought in for emergency surgery. In the quietest periods, the percentage can drop to 25 percent.

Older people often have to undergo surgery for hip [fractures](#). The average age of hip surgery patients in the study was 85, with women making up 72 percent. The median wait time before being operated on was about 20 hours.

Older people naturally have a greater risk of dying than the [average population](#) has. Age is one reason why mortality is high following surgery.

The capacity of the hospitals also plays a major role.

Previous results show that patients who are operated on for hip fractures have a higher risk of dying if they are discharged from the hospital early due to lack of space.

Hip fracture patients may need to be prioritized in the queue to increase their chances of survival.

Prioritizing patients "is part of a discussion about organizing emergency surgery. This could mean that we need to screen surgery [hip fracture](#) patients more than what's currently being done," says orthopaedist Lars Gunnar Johnsen.

More information: Sara Marie Nilsen et al, High volumes of recent surgical admissions, time to surgery, and 60-day mortality, *The Bone & Joint Journal* (2021). [DOI: 10.1302/0301-620X.103B2.BJJ-2020-1581.R1](#)

Provided by Norwegian University of Science and Technology

Citation: Hip fracture outcomes worse during busy periods (2021, March 10) retrieved 26 April 2024 from <https://medicalxpress.com/news/2021-03-hip-fracture-outcomes-worse-busy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.