

# One minute diagnostic found superior to standard tests for *P. vivax* malaria

March 15 2021

---



Gazelle is a compact, rugged, battery-operated diagnostic device. Credit: Hemex Health

Results of a clinical study conducted by researchers in Manaus, Brazil, show that the Gazelle Malaria test outperformed Rapid Diagnostic Tests

(RDTs) and was nearly as accurate as more expensive and time-consuming expert microscopy in detecting *Plasmodium vivax* (*P. vivax*) malaria. The data was published Friday, March 12 in the *Malaria Journal*.

The double-blind study was based on 300 participants who sought care from Fundação de Medicina Tropical Doutor Heitor Vieira Dourado (FMT-HVD), located in Manaus, an area in the Western Brazilian Amazon, heavily endemic for *P. vivax*.

"Because *P. vivax* is not adequately detected by current tests, the infection is frequently missed, and patients don't receive essential treatment," said Dr. Marcus Lacerda, Infectious Disease Researcher at FMT-HVD and lead investigator for the study.

"This causes recurrent symptoms and contributes to the spread of the disease. These results show that we're on a path to reduced suffering and faster malaria elimination."

*P. vivax* is the second most prevalent malaria species infecting humans and is widespread among many countries seeking to eliminate malaria. Although considered less deadly than *P. falciparum* (the most [common species](#)), *P. vivax* infections can result in serious illness and mortality.

Additionally, *P. vivax* presents a special challenge for diagnostics due to its typically low levels of parasitemia. Current RDTs miss many *P. vivax* infections, thus light microscopy, which is time-consuming and requires skilled technicians, is the current best option for many regions of the world.

Researchers at FMT-HVD noted that the Gazelle Malaria Test's use of hemozoin, a highly specific biomarker present in all species of [malaria](#), shows promise for detecting low parasitemia *P. vivax* infections.

When compared to optical microscopy in this study, the sensitivity and specificity of the Gazelle test were 96.2% and 100% respectively, whereas for RDTs they were 83.9% and 100%. This equates to RDTs missing 16 cases of *P. vivax* infection per 100 people, and Gazelle missing only 4.

The researchers also noted that Gazelle's portability, all-day battery operation, and supplies that do not require cold chain, make the device a promising alternative to [light microscopy](#) in field conditions.

"This milestone aligns with Hemex's strategy to assist countries with elimination efforts," said Patti White, CEO of Hemex Health. "We fully expect that with Gazelle's affordability, ease-of-use, and ultra-fast results, it can be used successfully for case management, mass and border screenings or during outbreaks."

The company has regulatory approval in a growing number of countries, including India, Kenya, and Ghana, and plans to apply for approval in South America soon.

**More information:** Gisely Cardoso de Melo et al, Performance of a sensitive haemozoin-based malaria diagnostic test validated for vivax malaria diagnosis in Brazilian Amazon, *Malaria Journal* (2021). [DOI: 10.1186/s12936-021-03688-0](#)

Provided by Hemex Health

Citation: One minute diagnostic found superior to standard tests for *P. vivax* malaria (2021, March 15) retrieved 5 May 2024 from <https://medicalxpress.com/news/2021-03-minute-diagnostic-superior-standard-p.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.