

Adversity in early life linked to higher risk of mental health problems

April 22 2021



Credit: CC0 Public Domain

New research has found that childhood adversity, such as parental conflict, death of a close family member or serious injury, before the age of nine was associated with mental health problems in late

adolescence.

However, the research also shows that improving the relationship between parents and children could prevent subsequent mental health problems, even in children who have experienced severe adversities. The research also indicated that improving a child's self-esteem and increasing their levels of physical activity can help to reduce the risk of developing mental health problems.

The study, led by researchers from RCSI University of Medicine and Health Sciences, is recently published in *Psychological Medicine*.

The research team analyzed data from over 6,000 children in Ireland who took part in the Growing Up in Ireland study. The results showed that just over a quarter of children had experienced childhood adversity before the age of nine.

At age 17 and 18, almost one in five of the [young people](#) were experiencing significant mental health difficulties. 15.2% had developed internalizing problems, such as anxiety or depression, and 7.5% had developed externalizing problems, such as conduct problems or hyperactivity.

Those who experienced childhood adversity were significantly more likely to report mental health problems in late adolescence.

Parent-child conflict explained 35% of the relationship between childhood adversity and late adolescent externalizing problems. The conflict also accounted for 42% of the relationship between childhood adversity and internalizing problems.

The child's self-esteem (also called [self-concept](#)) explained 27% of the relationship between child adversity and later internalizing problems.

The child's level of physical activity explained 9% of the relationship between [childhood adversity](#) and later internalizing problems.

"Children who experience multiple or severe life events are at an increased risk of mental health problems, but not all of those exposed to such events develop such problems. Our research points to some factors that can be useful for off-setting the risk of mental health problems in those who have been exposed to difficult life events," said Dr. Colm Healy, the study's lead author and postdoctoral researcher at RCSI University of Medicine and Health Sciences.

The work was funded by the Health Research Board in Ireland and the European Research Council.

"Among children who have experienced adversity, we found that reducing conflict between the parent and child and fostering a warm relationship can protect them from a broad range of later [mental health problems](#)," said Professor Mary Cannon, the study's principal investigator and professor of Psychiatric Epidemiology and Youth Mental Health at RCSI.

"We also found that improving a child's self-esteem and encouraging [physical activity](#) may also be useful intervention targets for preventing difficulties with mood and anxiety following earlier adversity. On the whole, this is a hopeful story that points towards effective interventions to improve outcomes for [children](#) who had experienced difficulties early in life."

More information: Colm Healy et al. Mediators of the longitudinal relationship between childhood adversity and late adolescent psychopathology, *Psychological Medicine* (2021). [DOI: 10.1017/S0033291721000477](#)

Provided by RCSI

Citation: Adversity in early life linked to higher risk of mental health problems (2021, April 22) retrieved 20 March 2024 from <https://medicalxpress.com/news/2021-04-adversity-early-life-linked-higher.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.