

To combat gum disease, help oral bacteria evolve

April 12 2021



Credit: Jonathan Borba from Pexels

Liver disease, from metabolic and bacterial causes, is a growing concern. What connects these dots? The gut, or more specifically, bacteria in the gut. Bacteria that cause inflammation in the mouth are transported



through the digestive tract to the gut and liver, where they can cause liver inflammation. Lipopolysaccharides, important structural molecules in some bacteria, act as endotoxins, producing systemic effects that can manifest as non-alcoholic fatty liver disease (NAFLD). Now, a multidisciplinary team from the University of Tsukuba show that exercise could be used to improve the oral environment in people with NAFLD, potentially leading to a new treatment for the disease.

These researchers previously demonstrated that exercise benefits patients with NAFLD by reducing fat, inflammation, and scarring in the liver; improving the liver's response to and clearance of the endotoxin; and reducing gum disease. With the latest study in their series, they add another signpost to uncharted territory:

"We know that exercise has innumerable benefits to health overall and for these specific conditions," says corresponding author Professor Junichi Shoda. "With this study, we sought to characterize underlying mechanisms—that is, show how exercise alters physiology and how altered physiology induces changes in oral <u>bacteria</u>."

The researchers carried out biochemical and genetic analyses on saliva from overweight men with NAFLD and gum disease before and after 12-week exercise or diet programs. Men in both groups were able to lose fat mass, but those following dietary restrictions also lost muscle mass, whereas those following the exercise program gained muscle mass. "More importantly, we found that reductions in lactoferrin, lipopolysaccharide, and IgA concentrations were only evident in the men who followed the exercise regimen," Professor Shoda explains, "which suggested that the oral environment had been significantly altered by exercise."

The samples from the exercise group also showed increased bacterial diversity and changes in the relative constituent bacterial populations. In



the overall population, more bacteria expressed genes related to environmental information processing, and less bacteria expressed genes related to genetic information processing and metabolism. In fact, bacteria expressed fewer genes for producing lipopolysaccharides.

"Therefore, it seems that, in people with both non-alcoholic <u>liver disease</u> and gum disease, <u>exercise</u> causes a biochemical shift in the environment of the mouth that favors the survival of less harmful bacteria," explains Professor Shoda.

A greater understanding of how to reduce the population or harmful effects of certain bacteria in patients with liver and periodontal disease could potentially lead to the development of new and improved treatment strategies for these diseases and other conditions thought to be influenced by bacteria, such as inflammatory bowel <u>disease</u>.

More information: Fumihiko Uchida et al. Effects of Exercise on the Oral Microbiota and Saliva of Patients with Non-Alcoholic Fatty Liver Disease, *International Journal of Environmental Research and Public Health* (2021). DOI: 10.3390/ijerph18073470

Provided by University of Tsukuba

Citation: To combat gum disease, help oral bacteria evolve (2021, April 12) retrieved 24 May 2024 from <u>https://medicalxpress.com/news/2021-04-combat-gum-disease-oral-bacteria.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.