

Discontinuing antidepressants requires careful planning

April 20 2021, by Riitta Partanen



Credit: Unsplash/CC0 Public Domain

The decision to stop prescribing long-term anti-depressants should be made only after careful preparation by GPs and their patients, University of Queensland research has found.

The [qualitative study](#) was based on interviews with 22 GPs working in urban and regional areas, who said there was no standardized approach.

Associate Professor Riitta Partanen from UQ's Rural Clinical School said GPs told researchers the discontinuation of such prescriptions was usually challenging.

"It begins with considering a patient's social and relational context, and is a journey involving careful preparation, tailored care and regular review," Dr. Partanen said.

"Stopping unnecessary long-term use needs to start at initiation, and GP education and training must give adequate attention to prescribing, repeat prescribing and deprescribing.

"The study suggests GPs need the clinical time to engage with patients with [decision](#)-making around prescribing and discontinuing antidepressants."

She said the research added to existing evidence that repeat [prescriptions](#) occurred within a complex medical and relational setting.

"They're never just administrative encounters to get a script," Dr. Partanen said.

"The study captures a set of steps GPs take together with their [patients](#) to initiate and set-up adequate support before, during and after discontinuation.

"It shows the complex decision-making GPs undertake in determining whether a patient is ready to discontinue."

According to the OECD, [Australians were the second largest per-capita](#)

[users of antidepressants](#) of the 30 countries for which data was available in 2000 and 2015.

Dr. Partanen said previous work had shown that long-term antidepressant use was on the rise and most antidepressant prescription took place in general practice.

"About one third of long-term users have no clear reason for continued use," she said.

"So there is scope within [primary care](#) for careful treatment re-evaluation and discontinuation of antidepressants."

The study is published in the *British Journal of General Practice*.

More information: Long-term antidepressant use in general practice: a qualitative study of GPs' views on discontinuation. *British Journal of General Practice*. DOI: doi.org/10.3399/BJGP.2020.0913

Provided by University of Queensland

Citation: Discontinuing antidepressants requires careful planning (2021, April 20) retrieved 20 March 2024 from <https://medicalxpress.com/news/2021-04-discontinuing-antidepressants-requires.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--