


# Good dental health may help prevent heart infection from mouth bacteria

April 15 2021

---



Credit: Pixabay/CC0 Public Domain

Maintenance of good oral health is more important than use of antibiotics in dental procedures for some heart patients to prevent a heart infection caused by bacteria around the teeth, according to a new

American Heart Association (AHA) scientific statement published today in the association's flagship journal, *Circulation*.

Infective endocarditis (IE), also called bacterial endocarditis, is a heart infection caused by bacteria that enter the bloodstream and settle in the heart lining, a heart valve or a blood vessel. It is uncommon, but people with heart valve disease or previous valve surgery, [congenital heart disease](#) or recurrent infective endocarditis have a greater risk of complications if they develop IE. Intravenous drug use also increases risk for IE. Viridans group streptococcal infective endocarditis (VGS IE) is caused by bacteria that collect in plaque on the tooth surface and cause inflammation and swelling of the gums. There's been concern that certain dental procedures may increase the risk of developing VGS IE in [vulnerable patients](#).

The new guidance affirms previous recommendations that only four categories of [heart patients](#) should be prescribed antibiotics prior to certain dental procedures to prevent VGS IE due to their higher risk for complications from the infection:

- those with prosthetic heart valves or prosthetic material used for valve repair;
- those who have had a previous case of infective endocarditis;
- adults and children with congenital heart disease; or
- people who have undergone a [heart](#) transplant.

"Scientific data since the 2007 AHA guidelines support the view that limited use of preventive antibiotics for dental procedures hasn't increased cases of endocarditis and is an important step at combating antibiotic overuse in the population," said Walter R. Wilson, M.D., chair of the statement writing group and a consultant for the Division of Infectious Diseases, Department of Internal Medicine at Mayo Clinic in Rochester, Minn.

It has been over a decade since recommendations for preventing infective endocarditis were updated amid concerns of antibiotic resistance due to overprescribing. The American Heart Association's 2007 guidelines, which presented the biggest shift in recommendations from the Association on the prevention of [infective endocarditis](#) in more than 50 years, more tightly defined which patients should receive preventive antibiotics before certain dental procedures to the four high-risk categories. This change resulted in about 90% fewer patients requiring antibiotics.

The scientific statement writing group reviewed data on VGS IE since the 2007 guidelines to determine if the guidelines had been accepted and followed, whether cases of and mortality due to VGS IE have increased or decreased, and if the guidance might need to be adjusted.

The writing committee reports their extensive review of related research found:

- There was good general awareness of the changes in the 2007 guidelines, however, adherence to the guidelines was variable. There was about a 20% overall reduction in prescribing preventive antibiotics among high-risk patients, a 64% decrease among moderate-risk patients, and a 52% decrease in those patients at low- or unknown-risk.
- In a survey of 5,500 dentists in the U.S., 70% reported prescribing preventive antibiotics to patients even though the guidelines no longer recommend it, and this was most often for patients with mitral valve prolapse and five other cardiac conditions. The dentists reported that about 60% of the time the antibiotic regimen was recommended by the patient's physician, and 1/3 of the time was according to patient preference.
- Since the stricter 2007 antibiotic guidelines, there is no convincing evidence of an increase in cases of VGS IE or

increased mortality due to VGS IE.

- The writing group supports the 2007 recommendation that only the highest risk groups of patients receive antibiotics prior to certain dental procedures to help prevent VGS IE.
- In the presence of poor oral hygiene and gingival disease, VGS IE is far more likely to develop from bacteria attributable to routine daily activities such as toothbrushing than from a dental procedure.
- Maintenance of good oral hygiene and regular access to dental care are considered as important in preventing VGS IE as taking antibiotics before certain dental procedures.
- It is important to connect patients with services to facilitate access to dental care and assistance with insurance for dental coverage, especially in those patients at high risk for VGS IE.
- It is still appropriate to follow the recommendation to use preventive [antibiotics](#) with high-risk patients undergoing [dental procedures](#) that involve manipulation of the gum tissue or infected areas of the teeth, or perforation of the membrane lining the mouth.

The scientific statement was prepared by the volunteer writing committee on behalf of the American Heart Association's Young Hearts Rheumatic Fever, Endocarditis and Kawasaki Disease Committee; the Council on Lifelong Congenital Heart Disease and Heart Health in the Young; the Council on Cardiovascular and Stroke Nursing; and the Council on Quality of Care and Outcomes Research.

**More information:** *Circulation* (2021). [DOI: 10.1161/CIR.0000000000000969](#)

Provided by American Heart Association

Citation: Good dental health may help prevent heart infection from mouth bacteria (2021, April 15) retrieved 5 May 2024 from <https://medicalxpress.com/news/2021-04-good-dental-health-heart-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.