

# More kids are being diagnosed with ADHD for borderline behaviors: Why that's a concern

April 13 2021, by Luise Kazda

---



Credit: CC0 Public Domain

During my daughter's challenging first year of school, we discovered how much effort it took her to sit and learn.

She was the youngest in her class, placing her at [higher risk](#) of being diagnosed with [ADHD](#) (attention deficit hyperactivity disorder).

While she struggled with attention and hyperactivity, her problems were always more frustrating than truly impairing. Still, constant battles over finishing tasks, the amount of time (and nerves) spent on a child that needs that extra bit of attention and the anger or sadness on her face made me wonder if we should try to get some support.

Maybe a diagnosis could be a straightforward fix to the problem?

## **What's the problem?**

Increasing awareness of ADHD has led to [consistent rises](#) in the number of [children](#) diagnosed with and [treated for](#) it, both internationally and in Australia. This would be good if it meant we were getting better at finding, diagnosing and helping children impaired by inattention or hyperactivity.

However, my [newly published study](#) in *JAMA Network Open* finds these increases in ADHD diagnoses may be largely due to children like my daughter, whose behaviors fall within a normal (but frustrating) range. I conducted this research with colleagues from the University of Sydney and Bond University.

Our study concluded these children are unlikely to benefit from being labeled with ADHD and may, in fact, be harmed by it.

This surge in diagnoses also results in limited resources being stretched thinner [among more children](#), ultimately taking away from those with severe problems who would benefit from more support.

## What is ADHD? And why is it so controversial?

ADHD [is a](#) "persistent pattern of inattention and/or hyperactivity-impulsivity that interferes with functioning or development".

It's one of the most common childhood disorders, affecting about [5-7%](#) of children. Over the past decades, debate on the appropriateness of diagnoses has grown in line with the rate of diagnosis.

Allen Frances, a prominent American psychiatrist, has been one of the most vocal critics of the trend. He [describes it](#) as the medicalisation of "every day experiences that are part of the human condition".

However, others suggest the increases in diagnosed children are largely due to [improved detection](#) in previously undiagnosed children.

[Both sides](#) of the debate claim to have proof. But we were surprised to discover no-one had ever summarized the scientific evidence for the key reasons behind increasing diagnosis rates.

So we reviewed the results from over 300 studies on ADHD over the past 40 years to determine which children are being newly diagnosed and if they benefit. Our [study design](#) allowed us to summarize a huge variety of studies in a way not done before.

## What we did and what we found

We found that since the 1980s, increasing numbers of school-aged children and adolescents around the world have been diagnosed with ADHD and medicated for it.

We know ADHD-related behaviors exist on [a spectrum](#) with no or

minimal hyperactivity and inattention on one end and severe ADHD on the other.

Many children can get distracted easily, are forgetful, find it difficult to sit still or wait their turn. In most children, these behaviors are mild enough to not interfere with a "normal" life.

However, there is no clear biological cut-off point above which someone just "has" ADHD. Ways of diagnosing ADHD also vary between countries and change over time, with criteria generally becoming less stringent.

Together, this ensures many potentially new cases could be discovered, depending on how low the bar is set.

[In the US](#), for example, almost half of all children diagnosed with ADHD have [mild symptoms](#), with only around 15% presenting with severe problems. Only about [1% of all children](#) in an Italian study had severe ADHD-related behaviors. And, in general, children today are [no more hyperactive or inattentive](#) than 20 years ago.

All this led us to conclude a substantial proportion of these additional diagnoses (children who would not have been diagnosed 20 years ago) are, at best, borderline cases.

For example, [one study](#) shows while diagnoses increased more than five-fold over ten years in Sweden, there was no increase in clinical ADHD symptoms over the same time. This means that with the lowering of the diagnostic bar, children diagnosed with ADHD are, on average, [less impaired](#) and more similar to those without an ADHD diagnosis.

As a result children like my daughter, who are [the youngest in their class](#), are at risk of being labeled with ADHD because their relative

immaturity can be enough to push them over the threshold into the zone of "abnormal" behavior.

## Why it's important to get it right

### For children with mild symptoms

Children with mild ADHD symptoms are unlikely to benefit from a diagnosis. They (and their families) also incur [substantial costs](#) as well as [potential harms](#) from the diagnosis and treatment. That's because:

- instead of drumming up extra support, an ADHD label can have [negative](#) social, psychological and academic effects, when compared to similar [young people](#) without a diagnosis
- medication [reduces symptoms to a lesser extent](#) in children with mild ADHD (however it is beneficial in many severe cases)
- medication for young people with milder symptoms also has no positive, but a potential negative, effect on [academic outcomes](#) (such as maths and reading scores) when compared to unmedicated young people with similar behavior. Also, medication doesn't reduce the risks of [injuries, criminal behavior](#) and [social impairment](#) as much as in those with severe symptoms.

### For children with severe symptoms

It's also important children with more severe ADHD symptoms are correctly diagnosed so they don't miss out on much-needed support.

With [ever-increasing diagnosis rates](#) of ADHD, schools are increasingly struggling to adequately support every child with a diagnosis: the slice of funding and support every child can receive gets smaller and smaller, the more children are included.

In turn, this often means those with the most severe problems [get left behind](#).

## What can we do?

In light of the potential risks associated with diagnosing a [child](#) with milder ADHD symptoms, we recommend doctors, parents and teachers work together following a "[stepped diagnosis approach](#)". This ensures swift and efficient [diagnosis](#) and treatment in severe cases. For those with milder symptoms, taking some time to watch and wait may mean many of them won't need to be labeled or treated.

Not only will this avoid potential harm for individual children, it also ensures resources are allocated where they are needed most and will be most effective.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: More kids are being diagnosed with ADHD for borderline behaviors: Why that's a concern (2021, April 13) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-04-kids-adhd-borderline-behaviors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--