

# Minimally invasive retinal reattachment procedure leads to superior photoreceptor integrity

April 22 2021

---



Credit: Unsplash/CC0 Public Domain

A minimally invasive retinal reattachment procedure that can be done in an ophthalmologist's office leads to better long-term integrity and

structure of the retina's photoreceptors—cells that allow us to see—compared with more invasive operating room procedures, according to new research published April 22.

The study, published in *JAMA Ophthalmology* and led by researchers at St. Michael's Hospital of Unity Health Toronto, contributes to a growing body of evidence pointing towards pneumatic retinopexy (PnR) as the better first-line retinal reattachment technique to achieve the best visual outcomes.

Retinal detachment is the most common surgical ocular emergency, progressing to loss of vision within hours or days. While there are a number of different treatment options, there is limited high-quality data guiding the choice of surgery.

"Our study shows there is a difference in the long-term integrity of the photoreceptors between different surgical techniques, and these anatomic differences were associated with visual outcomes," said co-principal investigator Dr. Rajeev H. Muni, a vitreoretinal surgeon at St. Michael's and researcher at the Li Ka Shing Knowledge Institute.

Previous studies carried out at St. Michael's Hospital demonstrated that patients had better visual results following the less invasive PnR compared to pars plana vitrectomy (PPV), an alternative operating room procedure. In this study, the researchers determined there was an actual difference in photoreceptor anatomic recovery between the two retinal reattachment techniques.

Using data from a randomized trial conducted at St. Michael's Hospital, researchers compared the retinal scans of 72 patients who had retinal reattachment using PPV and 73 patients who had retinal reattachment using the minimally invasive PnR at 12 months post-operatively.

PnR is a less invasive and less expensive retinal reattachment technique that can be done in an ophthalmologist's office. In PnR, a small gas bubble is used to close the retinal tear and allow the fluid to reabsorb naturally and slowly. PPV is a surgical technique where the fluid under the retina is rapidly aspirated and removed, forcefully bringing the retina back in position.

The imaging showed that discontinuity—an absence of a part of the photoreceptor layer—was more common at 12 months post-operatively among patients who had the PPV surgical reattachment than in patients who had the minimally invasive PnR procedure. Discontinuity indicates damage to cells that are critical for vision and this damage was found to be associated with worse visual outcomes.

"This data provides an objectively determined anatomic basis for the superior functional outcomes seen with pneumatic retinopexy that we have previously reported," said Dr. Roxane J. Hillier, the trial's other co-principal investigator who is now based in the United Kingdom.

The researchers say their findings highlight that closing the retinal tear, doing as little else as possible, and allowing the retina to reattach naturally leads to the best outcomes both from an anatomic and visual perspective.

"This was previously unknown, and in my opinion will be a game-changer in our field," said Dr. Muni.

**More information:** Rajeev H. Muni et al. Postoperative Photoreceptor Integrity Following Pneumatic Retinopathy vs Pars Plana Vitrectomy for Retinal Detachment Repair: A Post Hoc Optical Coherence Tomography Analysis From the Pneumatic Retinopexy Versus Vitrectomy for the Management of Primary Rhegmatogenous Retinal Detachment Outcomes Randomized Trial. *JAMA Ophthalmol*. Published online April

22, 2021. [DOI: 10.1001/jamaophthalmol.2021.0803](https://doi.org/10.1001/jamaophthalmol.2021.0803)

Provided by St. Michael's Hospital

Citation: Minimally invasive retinal reattachment procedure leads to superior photoreceptor integrity (2021, April 22) retrieved 3 May 2024 from <https://medicalxpress.com/news/2021-04-minimally-invasive-retinal-reattachment-procedure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.