

# Neuro-researchers find repetitive head impacts can result in functional brain impairments

April 22 2021

---



Hockey player gets his brain vital signs checked. Researchers find repetitive head impacts can result in functional brain impairments in youth hockey. Credit: Health and Technology District

Neuroscience researchers at Mayo Clinic Orthopedics and Sports Medicine in Rochester, Minnesota, U.S., the Health and Technology District and Simon Fraser University (SFU) in Surrey, British Columbia, Canada have published the latest results of their ongoing multi-year hockey concussion study examining changes in subconcussive cognitive brain function in male youth ice hockey players.

The research team monitored [brain vital signs](#) during pre- and post-season play in 23 Bantam (age 14 or under) and Junior A (age 16 to 20) male ice-hockey players in Rochester, Minnesota.

"Brain vital signs" translates complex brain waves measured using portable electroencephalography (EEG) at the rink-side, into simple, fast, user-friendly and intuitive results that provide an objective evaluation of cognitive brain function. Called the ABCs of brain function, brain vital signs track three well-established neural responses for Auditory sensation (using a response called the N100), Basic attention (using a response called the P300), and Cognitive processing (using a response called the N400).

The study builds on 2019 results published in *Brain: A Journal of Neurology* that confirmed significant brain vitals sign changes shortly after concussions were diagnosed in Junior A players. Key results of this study showed undetected impairments remained when players were cleared to return to play using current clinical concussion protocols. Notably, the initial study also reported sensitivity to subconcussive impairments in those players who did not sustain a concussion diagnosis over the course of the season.

The current second phase of the study replicated these results and added the Bantam age group. The latest findings were recently published in the peer-reviewed journal *Brain Communications* in the advanced articles section.

The results of this new *Brain Communications* study showed:

- Significant brain vital sign changes in N100 Auditory sensation and N400 Cognitive processing responses for the pre-to-post season comparison across both groups.
- Differences between the Bantam and Junior A ice hockey players showed more changes in the Junior A group.
- Importantly, the subconcussive changes were significantly correlated with the number of head impacts over the season across both age groups and consequently showed more subconcussive changes in brain vital signs.

A subconcussive impact is a mechanical force transmitted to the brain below the threshold for a diagnosis of an acute concussive injury. The effects of these low-magnitude impacts may not even be noticeable to the player or to observers on the sideline. Head impacts in the sport of ice hockey typically result from player-to-player or player-to-boards contact due to body checking, collisions and fighting . Some of these impacts are the consequence of foul play, but many of these events also result from routine, legal on-ice behavior.

"Concussion in sports is a major concern for many and our research has shown that having an objective physiological measure of brain function at rink-side is key to detection and managing concussive impacts," says Dr. Aynsley Smith, principal investigator of the study and Associate Professor of Orthopedics at Mayo Clinic in Rochester, Minnesota.

The study, funded in part by USA Hockey, was co-led by Dr. Michael Stuart, Professor of Orthopedics at Mayo Clinic in Rochester, Minnesota. It is also part of a larger concussion research team within Mayo Clinic that includes neurology research leadership from Dr. David Dodick, Professor of Neurology at Mayo Clinic in Phoenix, Arizona, U.S.

Canadian collaborators include neuroscientist Dr. Ryan D'Arcy, an SFU professor and co-founder of the Health and Technology District in Surrey B.C. and Dr. Shaun Fickling, a biomedical engineer, a recent SFU Ph.D. graduate and lead author of the study.

"Our research has shown that repetitive subconcussive impacts triggered compounding effects in brain function changes, which underscores the importance of shifting our thinking and understanding of concussions as a singular acute-injury model to a spectrum of head-impact exposure and effects over time," says Dr. Fickling.

The US-Canadian concussion research team is continuing to advance their collaborative effort.

Says Dr. D'Arcy, "In medicine: you can't treat what you can't measure. With breakthroughs on measurement challenges, we hope to now accelerate treatment innovations for prevention, acute care and extended care concussion management—for all people across a range of different applications. Our partnership is moving into incredibly exciting future steps—stay tuned."

**More information:** Shaun D Fickling et al, Subconcussive brain vital signs changes predict head-impact exposure in ice hockey players, *Brain Communications* (2021). [DOI: 10.1093/braincomms/fcab019](https://doi.org/10.1093/braincomms/fcab019)

Provided by Simon Fraser University

Citation: Neuro-researchers find repetitive head impacts can result in functional brain impairments (2021, April 22) retrieved 18 April 2024 from <https://medicalxpress.com/news/2021-04-neuro-researchers-repetitive-impacts-result-functional.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.