

# During pandemic's first 9 months, depression and anxiety increased sharply among Americans

April 22 2021

---



Credit: Pixabay/CC0 Public Domain

Confirming anecdotal evidence that the spread of the coronavirus has strained Americans' mental health, Boston College researchers found

reports of anxiety increased to 50 percent and depression to 44 percent by November, 2020—rates six times higher than 2019—according to a new report in the journal *Translational Behavioral Medicine*.

Among U.S. adults aged 18-29, the impact on mental [health](#) was even more severe. Rates of anxiety and depression increased to 65 percent and 61 percent, respectively, of the respondents in that age group, according to the report.

Use of prescription medication, counseling services, and unmet need for mental health services also rose significantly, according to the co-authors of the new study, Boston College developmental psychologist Rebekah Levine Coley and economist Christopher F. Baum, who reviewed [survey data](#) from nearly 1.5 million U.S. adults.

Rates of mental health disorders were highest among young, less-educated, single parent, female, Black and Hispanic respondents, Coley and Baum report. Disparities between young versus older and less educated versus more educated adults rose over time. Young, female, and moderately educated respondents also reported higher unmet needs for services.

"Disparities in estimates of mental health disorders and mental health treatment indicate a striking disequilibrium between the potential need for and the use of mental health services during the COVID-19 pandemic," said Coley, a professor in the Lynch School of Education and Human Development. "Rising mental health challenges are being borne largely by young, less advantaged people of color, and women, with the potential for expanded interruptions to optimal functioning and societal recovery from COVID-19."

Despite extensive anecdotal evidence of rising mental health challenges posed by the pandemic, little prior evidence had systematically assessed

rates of mental health disorders or use of mental health symptoms through the course of the COVID-19 pandemic, said Coley.

"We discerned a need to track rates of depression and anxiety, as well as rates of use of mental health services and reports of unmet need for such services between April and November, 2020," she said. "We also sought to assess whether the rates of mental health disorders and [service](#) use varied across key demographic groups in the U.S."

The researchers analyzed data gathered between April and November 2020 as part of the U.S. Census' Household Pulse Survey, a series of cross-sectional surveys conducted weekly. Survey respondents self-reported their symptoms of anxiety and depression, use of medication, use of counseling services, and unmet need for services. Coley and Baum analyzed these surveys to track trends in mental health disorder symptoms and access to and use of mental health services.

While the researchers expected to see increases in the rates of depression and anxiety, they were surprised by the magnitude of the increases, Coley said.

"The fact that prevalence rates were six times higher than national norms from 2019 was striking, as was the fact that these increases were born primarily by younger adults, aged 18-29 years, whose rates of anxiety and depression were nearly twice as high as those of older adults, aged 70 and above," Coley said.

The analysis reveals a need for continued study of the pandemic's impact on mental health.

The findings suggest the need for increased access to [mental health services](#) and other supports to help adults face the economic, social, and psychological stressors of the COVID-19 pandemic, Coley said. There is

also a need to assess whether rising mental health challenges continue to grow as the pandemic rages on, and to delineate the longer-term effects of the social, economic, and psychological disruptions caused by COVID-19. For [young adults](#) particularly, the pandemic may have interrupted higher education plans and the initiation of careers and social and financial independence, with unknown long-term repercussions.

"The myriad stressors imposed by the COVID-19 pandemic have impaired mental health and well-being," said Baum, who also holds an appointment at BC's School of Social Work. "Although evidence from early in the pandemic revealed elevated rates of [mental health](#) conditions, research had not documented whether psychological disorders have continued to rise as the [pandemic](#) has persisted, or for whom they have risen most dramatically."

**More information:** Rebekah Levine Coley et al, Trends in mental health symptoms, service use, and unmet need for services among U.S. adults through the first 9 months of the COVID-19 pandemic, *Translational Behavioral Medicine* (2021). [DOI: 10.1093/tbm/ibab030](https://doi.org/10.1093/tbm/ibab030)

Provided by Boston College

Citation: During pandemic's first 9 months, depression and anxiety increased sharply among Americans (2021, April 22) retrieved 28 April 2024 from <https://medicalxpress.com/news/2021-04-pandemic-months-depression-anxiety-sharply.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---