

## Study examines racial inequity in suicide prediction models

April 28 2021



Credit: Unsplash/CC0 Public Domain

Models that can successfully predict suicides in a general population sample can perform poorly in some racial or ethnic groups, according to a study by Kaiser Permanente researchers published April 28 in *JAMA* 



## Psychiatry.

The new findings show that 2 <u>suicide</u> prediction models are less accurate for Black, American Indian, and Alaska Native people and demonstrate how all prediction models should be evaluated before they are used. The study is believed to be the first to look at how the latest statistical methods to assess suicide risk perform when tested specifically in different ethnic and <u>racial groups</u>.

More than 47,500 people died from suicide in the United States in 2019, an increase of 33% since 1999. Health care leaders hope to reduce suicide by using records of mental health visits, diagnoses, and other factors to identify patients at highest risk of suicide and intervene. In recent years, the Veterans Health Administration, HealthPartners, and several Kaiser Permanente regions have started using suicide prediction models to guide care.

"With enthusiasm growing for suicide prediction modeling, we must be sure that such efforts consider health equity," said Yates Coley, Ph.D., the study's first author and an assistant investigator at Kaiser Permanente Washington Health Research Institute. "Current methods maximize predictive performance across the entire population, with predictive accuracy in the largest subgroup—white patients—eclipsing the performance for less-prevalent racial and ethnic subgroups."

The *JAMA Psychiatry* article follows several studies that have uncovered similar concerns with racial/ethnic bias in algorithms in domains ranging from criminal justice and policing to health care. In the new study, Coley and colleagues gathered electronic health records for nearly 14 million outpatient mental health visits over a 7-year period from 7 health care systems.

Using these health records, the research team developed 2 different



models—a standard statistical logistic regression approach and a random forest machine learning algorithm—to predict suicide deaths within 90 days of a mental health visit. The models used demographic characteristics, comorbidities, mental health and substance use diagnoses, dispensed psychiatric medications, prior suicide attempts, prior mental health encounters, and responses to Patient Health Questionnaire 9, which is filled out routinely at mental health visits.

The models had a high degree of precision in identifying suicides and avoiding false positives across the entire sample, and for white, Hispanic, and Asian patients. The models fared much worse with Black, American Indian, and Alaska Native patients and patients without race/ethnicity recorded.

For example, after one of the models was used to predict the 5% of visits with the highest risk of suicide, the researchers zeroed in on white patient visits in that risk group and found that nearly half of eventual suicides were identified. By contrast, that risk group only included about 7% of suicide deaths by American Indian, Alaska Native, and Black patients. This means that people in these populations who die by suicide would be much less likely to be identified by the <u>model</u>, indicating that this particular tool is not useful for guiding their care.

- Researchers cited several possible reasons for shortcomings in prediction accuracy that could be the result of embedded biases in the data. Black, American Indian, and Alaska Native patients face barriers to accessing mental health services. This means that there's less data on <u>suicide risk</u> factors so it may be more difficult to make accurate forecasts.
- Even when these populations do have access to mental health services, research shows that they are less likely to be diagnosed and treated for mental health conditions. The clinical data may not accurately reflect risk, affecting the models' suicide



predictions.

• Suicides in these populations may be incorrectly identified as unintentional or accidental, contributing to the challenge in predicting suicides in these populations.

The 2 prediction models in the study are not the same as the ones now being implemented in health systems. This study examined models that predict suicide deaths, while models being used in clinical care at Kaiser Permanente predict self-harm, or suicide attempts. An audit, like the one in this study, of the suicide-attempt prediction model in use at Kaiser Permanente in Washington did not find racial or ethnic disparities in prediction performance. Such audits may be needed at other health organizations using suicide prediction models, and the study lays out the steps that should be taken when implementing a prediction model to ensure inequities are not perpetuated.

"Before we implement any prediction model in our clinics, we must test for disparities in accuracy and think about possible negative consequences," said Gregory Simon, MD, MPH, a study co-author and Kaiser Permanente Washington Health Research Institute senior investigator. "Some <u>prediction models</u> pass that test and some don't, but we only know by doing research like this."

Provided by Kaiser Permanente

Citation: Study examines racial inequity in suicide prediction models (2021, April 28) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2021-04-racial-inequity-suicide.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.