

Surgical technique may reduce positive margin rate in breast cancer patients with DCIS

April 5 2021



Credit: Pixabay/CC0 Public Domain

According to the American Cancer Society, a noninvasive breast cancer called ductal carcinoma in situ (DCIS) accounts for approximately one

of every four new breast cancer cases in the United States. If left untreated, DCIS has the potential to evolve into invasive cancer, so many patients choose to have breast-conserving surgery or mastectomy after a diagnosis.

However, obtaining clear or negative margins—no [cancer cells](#) in the outer edge of removed tissue—is critical to mastectomy success as positive margins are associated with higher rates of recurrence.

A new study from Wake Forest School of Medicine suggests removing additional tissue around the tumor cavity, a technique known as cavity shave margin resection, reduces the rate of positive margins in patients with DCIS. The findings are published online in the April 2021 issue of the *Journal of the American College of Surgeons*.

"Our previous clinical trial showed that this technique can reduce positive-margin rates in invasive disease by at least 50%, but to my knowledge, this is the first analysis involving pure DCIS," said Marissa Howard-McNatt, M.D., director of the Breast Cancer Center at Wake Forest Baptist Health and associate professor of surgical oncology at Wake Forest School of Medicine, part of Wake Forest Baptist.

Researchers randomized 109 patients with pure DCIS (no [invasive cancer](#)) from 10 centers across the U.S., and 58 (53.2%) of these patients were randomized for cavity shave margin resection. In this group, researchers noted a nearly 65% reduction in the positive-margin rate.

According to Howard-McNatt, achieving clear margins in patients with DCIS can be difficult due to the continuous and branching nature of the spread of this disease. Clear margins free of cancer are vital in decreasing the chance of recurrence of [cancer](#) in the [breast](#).

"Our study shows a significant reduction in positive margins," Howard-McNatt said. "This technique should be considered for DCIS patients."

More information: Marissa Howard-McNatt et al, Impact of Cavity Shave Margins on Margin Status in Patients with Pure Ductal Carcinoma In Situ, *Journal of the American College of Surgeons* (2020). [DOI: 10.1016/j.jamcollsurg.2020.11.019](https://doi.org/10.1016/j.jamcollsurg.2020.11.019)

Provided by Wake Forest University Baptist Medical Center

Citation: Surgical technique may reduce positive margin rate in breast cancer patients with DCIS (2021, April 5) retrieved 5 May 2024 from <https://medicalxpress.com/news/2021-04-surgical-technique-positive-margin-breast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.