

Two-color reporter shows cholinergic neurons engage ISR to modulate dopamine levels

April 26 2021, by Bob Yirka



Using a two-color viral reporter of ISR activation state, striatal CINs were distinguished from other cell types by showing steady-state ISR activation. ISR inhibition altered CIN excitability, inverted the sign of dopamine modulation of CIN firing rate and evoked dopamine release, and increased movement vigor in learned tasks. mAHP, medium afterhyperpolarization; AP, action potential; DA,



dopamine. Credit: Science (2021). DOI: 10.1126/science.abe1931

A team of researchers from Duke University Medical Center, the California Institute of Technology, the National Institute of Mental Health and Baylor College of Medicine has found via a new reporter that cholinergic neurons engage the integrated stress response (ISR) to modulate dopamine levels. In their paper published in the journal *Science*, the group describes their study of the ISR pathway using the new reporter and what they learned. Anna Ingebretson and Julia Lemos with the University of Minnesota have published a Perspectives piece in the same journal issue outlining current research involving striatal cholinergic interneurons and the work by the team on this new effort.

Prior research has shown that the <u>dorsal striatum</u>, situated in the <u>basal</u> ganglia, plays a role in planning, learning, movement, reward learning and the process of making and obtaining desired goals. The basal ganglia and its parts have been the target of much research in recent years because of its involvement in a wide range of activities and because it is believed problems with it are involved with a slew of major illnesses, including Parkinson's disease, substance abuse and depression. In this new effort, the researchers sought to learn more about the dorsal striatum and the ISR—a <u>signaling pathway</u> that prior research has shown plays an important role in problems with homeostasis in the basal ganglia.

To gain a better perspective on the role of ISR, the researchers developed a two-color reporter they called SPOTlight aimed at reporting on ISR activation states. Its use involved creating a viral vector to make different kinds of neurons respond differently to different types of light in mice. It was used to translate red or green fluorescent proteins produced due to signaling in the brain.



Prior research had shown that cholinergic interneurons make up less than 2% of the neurons in the dorsal striatum, yet they still play a major role in signaling in the ISR—such signaling has been shown to result in the production of the neurotransmitter acetylcholine. Using SPOTlight, the researchers were able to see that in mice, the cholinergic interneurons were also needed by the ISR to maintain properties related to neuronal firing and in regulating a response to the presence of dopamine.

More information: Ashley R. Helseth et al. Cholinergic neurons constitutively engage the ISR for dopamine modulation and skill learning in mice, *Science* (2021). DOI: 10.1126/science.abe1931

Anna E. Ingebretson et al. A spotlight on the elusive striatal cholinergic interneuron, *Science* (2021). <u>DOI: 10.1126/science.abi4907</u>

© 2021 Science X Network

Citation: Two-color reporter shows cholinergic neurons engage ISR to modulate dopamine levels (2021, April 26) retrieved 10 May 2024 from <u>https://medicalxpress.com/news/2021-04-two-color-cholinergic-neurons-engage-isr.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.