

# One bone fracture increases risk for subsequent breaks in postmenopausal women

May 5 2021

---



Credit: CC0 Public Domain

Current guidelines for managing osteoporosis specifically call out hip or spine fractures for increasing the risk for subsequent bone breaks. But a new UCLA-led study suggests that fractures in the arm, wrist, leg and other parts of the body should also set off alarm bells.

A fracture, no matter the location, indicates a general tendency to break a bone in the future at a different location, said Dr. Carolyn Crandall, the study's lead author and a professor of medicine at the David Geffen School of Medicine at UCLA.

"Current clinical guidelines have only been emphasizing hip and [spine fractures](#), but our findings challenge that viewpoint," Crandall said. "By not paying attention to which types of [fractures](#) increase the risk of future fractures, we are missing the opportunity to identify people at increased risk of future fracture and counsel them regarding risk reduction.

"Postmenopausal women and their physicians may not have been aware that even a knee fracture, for example, is associated with increased risk of future fractures at other locations of the body."

The study will be published May 5 in the peer-reviewed journal *EClinicalMedicine*.

The researchers examined records from 1993 through 2018 for more than 157,000 women aged 50 through 79. Data was sourced from the Women's Health Initiative, a national study funded by the National Heart, Lung, and Blood Institute.

The researchers found that among postmenopausal women, initial fractures of the lower arm or wrist, upper arm or shoulder, upper leg, knee, lower leg or ankle, and hip or pelvis were associated with an approximately three- to six-fold increase in risk for subsequent fractures. That finding held for all of the age groups studied, with higher risks being more pronounced among non-Hispanic Black, Hispanic or Latina, and Asian Pacific Islander women than among non-Hispanic White women.

The authors noted some limitations to the study, including the fact that the fractures were self-reported by participants. However, earlier research has demonstrated that statistics for self-reported fractures is fairly accurate compared with statistics from medical records.

Also, the researchers did not have information about broken ribs, which may have led them to underestimate the risk for other fractures—it's possible that the true effect could be even more pronounced than the results show—and bone mineral density was measured for only a subset of participants, so the researchers could not investigate whether the risk for future fractures was associated with bone density.

Although there is a need for more studies to understand why [women](#) of some ethnicities have a greater risk for a subsequent fracture following an initial bone break, the researchers write that their findings "indicate that aggressive follow-up of [postmenopausal women](#) who experience initial fracture is indicated. Our results will inform counseling, future guidelines, and the design of intervention trials regarding the selection of appropriate candidates for pharmacotherapy."

**More information:** *EClinicalMedicine*, [DOI: 10.1016/j.eclinm.2021.100826](#)

Provided by University of California, Los Angeles

Citation: One bone fracture increases risk for subsequent breaks in postmenopausal women (2021, May 5) retrieved 13 May 2024 from <https://medicalxpress.com/news/2021-05-bone-fracture-subsequent-postmenopausal-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.