

Concussions affect men and women differently: Female athletes need more attention in brain research

May 5 2021, by Shreya McLeod and James F. Donnelly



Credit: AI-generated image ([disclaimer](#))

News emerged last week that AFLW player [Jacinda Barclay](#), who died last year at age 29 following a short period of mental illness, had abnormalities in her brain tissue.

Barclay was the first Australian contact sportswoman to have her [brain](#) donated to the Australian Sports Brain Bank, a medical laboratory that investigates changes in [complex nerve structures](#) after death, in order to understand brain conditions sustained by sportspeople.

Similar to findings seen in some male athletes internationally, the researchers found changes in Barclay's white matter.

White matter has to do with neural connectivity in the brain. As white matter degrades, as is often seen in aging and in diseases such as Alzheimer's, [neural connectivity diminishes](#), contributing to cognitive decline.

However, we don't understand enough about this process in the brain—particularly in sportswomen.

Women's sport is becoming more popular

Over the past decade, Australia's elite women's sports have experienced unprecedented growth. More than [550,000 Australian women](#) (aged 14 and over) now play a form of football, be it soccer, Australian rules, rugby league or rugby union.

This increase in participation in contact and collision sports has seen a concurrent rise in injuries [such as concussions](#). Concussion is [a transient injury to the brain](#), caused by a jolt to the head or body.

Although specialized brain imaging can detect microscopic changes in the connections between brain regions, [concussion](#) is hard to detect on routine brain scans. Diagnosis typically relies on [people reporting symptoms](#) such as dizziness, confusion, unsteadiness, nausea and headaches after a collision.

Recognizing when someone has suffered a concussion, removing them from the field and carefully assessing recovery [are all crucial steps](#) before they return to play.

Risk factors for female athletes

Female athletes are [more likely](#) to sustain a concussion than their [male counterparts](#).

Like males, women report [a range of symptoms](#) after a concussion, such as headaches, mental fatigue, concentration difficulties and mood swings.

Although symptoms [can last longer](#) in some people, recovery from a concussion normally takes [seven to ten days](#) for adults. Research on length of recovery is mixed but overall supports that women [take longer to recover](#) than men.

Women also perform worse on [neurocognitive testing](#) post-injury, which measures things like decision-making ability and processing.

These gender-based differences may be due to a combination of factors.

Women tend to be more aware of their symptoms and are [more likely to report them](#), so this may account for some degree of the gap. However, under-reporting still exists.

Female athletes also generally have shorter and narrower necks, and lower head mass (their heads are smaller and less dense). These factors are associated with [lower neck strength](#). Neck strength is a [protective factor](#) against concussion, so women may be more susceptible for this reason.

Further, female brains metabolize glucose (sugar) and oxygen [faster than male brains](#). If a head injury temporarily disrupts blood supply to the brain, it could have a greater effect on the faster nutrient-burning female brain.

Meanwhile, sex hormones such as progesterone that [vary across a women's menstrual cycle](#) could also affect outcomes after a concussion. Concussions sustained during the follicular phase (after menstruation, before ovulation) are less likely to lead to symptoms a month later, whereas injuries in the luteal phase (after ovulation, before menstruation) result in [poorer outcomes](#).

Although we don't fully understand why, concussion outcomes appear to be worse when progesterone levels are high. However, these effects may be negated when women take the [contraceptive pill](#).

Despite these differences, women are [an understudied population](#) when it comes to concussion, resulting in a lack of gender-specific treatment guidelines.

Repeated concussions and long-term risk: a complex area

In the context of long-term brain injury and sport, we've perhaps most often heard about a condition called chronic traumatic encephalopathy (CTE).

Male former AFL players [Danny Frawley](#) and [Shane Tuck](#) were both diagnosed with CTE when their brains were examined after their deaths.

CTE is described as a delayed-onset and progressive neurodegenerative disease, with symptoms appearing in midlife or decades after exposure

to head traumas. It's believed these changes lead to an abnormal buildup of a protein called "tau," which can damage brain cells.

But CTE is not the only way in which changes to the brain might present over time. To date, no sportswoman has been diagnosed with CTE (including Barclay).

An article in [The Guardian](#) reported Barclay "did not have a substantial clinical history of concussion and her brain did not show evidence of her having sustained a concussion in the weeks before she died."

Brain bank research can detect white or gray matter changes but may not be able to ascertain which of several possible factors (for example, concussions, substance use, undiagnosed mental illness, aging) [led to the development](#) of brain pathology in a given case.

Brain changes seen in some deceased athletes have also been found in people with a variety of neurological and psychiatric disorders, but with [no known history](#) of head trauma or participation in risky activities such as contact sports or military combat.

So there's a lot we don't know, and more research we need to do, including involving people outside professional sport.

Increased focus on women is important

Women have [higher rates of Alzheimer's disease](#) than men. Although a history of head trauma [is a potential risk factor](#) for developing dementia and other neurodegenerative diseases like Parkinson's, studies have not yet examined [the interaction](#) between sport-related concussions and neurodegenerative diseases in [women](#).

We need [studies in women](#) to assess the interaction between exposure to

single or repeated head impacts and the [potential changes](#) in brain and behavior across the lifespan.

These studies need to use precise tests of attention, response speed and other cognitive abilities, and include indices of genetic [risk factors](#), mental health, and menstrual cycle function pre- and post-injury.

Barclay's groundbreaking donation to the Australian Sports Brain Bank is an important step towards gender equity in concussion research.

But to further advance our understanding, brain donors should participate in long-term studies during their lives that consider multiple causal or protective factors.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Concussions affect men and women differently: Female athletes need more attention in brain research (2021, May 5) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-05-concussions-affect-men-women-differently.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.