


# Engineering diseased human skin in vitro

May 10 2021

---



(A) Representative image showing diabetic foot ulcer caused by delayed wound healing in diabetic patients. (B) 3D cell printing system showing the materials and printing methods required for this study. (C) Modeling diabetic epidermis through intercellular interaction between diabetic human dermal fibroblasts and normal human epidermal keratinocytes, and a wounded skin model with delayed reepithelization. (D) Adding subcutaneous layer (blood vessel + fat) to better recapitulate pathophysiological functions of diabetes and to test applicability to drug screening platform.

Credit: Pohang University of Science & Technology (POSTECH)

The skin, which covers the surface of the human body, is its largest organ. It is the first organ to show changes stemming from organ or physiological activity. It is especially common for diabetic patients to suffer from skin diseases or infections. Recently, a POSTECH research team has succeeded in creating a 3D artificial skin that enables observation of skin diseases of diabetic patients.

A research team led by Professor Dong-Woo Cho and Minjun Ahn of POSTECH's Department of Mechanical Engineering and Professor Byoung Soo Kim of School of Biomedical Convergence Engineering at Pusan National University has successfully produced an in vitro diseased [skin](#) model that displays the pathophysiological hallmarks of type 2 diabetes based on a 3D cell printing system. These research findings were recently published in *Biomaterials*, a world-renowned international journal in biomaterials.

Despite continuous research to produce [artificial skin](#) with 3D cell printing technology, artificial skin displaying the pathological process present in the native skin has not been reported yet.

Inspired by the interaction between the epidermis and skin cells found in real skin, the research team hypothesized that when normal keratinocytes interact with the dermal layer made of diabetic fibroblasts, they will differentiate into diabetic epidermis. To prove this, diabetic artificial skin with skin wounds based on 3D printing technique was fabricated using each cell.

In this diabetic artificial skin, slow re-epithelialization, a typical feature of diabetic skin, was observed. In addition, when the diabetic fat tissue layer containing [blood vessels](#) was added, insulin resistance, adipocyte hypertrophy, pro-inflammatory response, and vascular dysfunction, which are commonly observed in diabetes, were confirmed.

"Through 3D cell printing, we can now observe [skin diseases](#) in vitro, without actually experiencing it," remarked the researchers. "We anticipate it to be a way to replace animal models that have been conventionally used to observe skin diseases. It is significant that its applicability as a disease model for new drug development has been proven."

**More information:** Byoung Soo Kim et al, Engineering of diseased human skin equivalent using 3D cell printing for representing pathophysiological hallmarks of type 2 diabetes in vitro, *Biomaterials* (2021). [DOI: 10.1016/j.biomaterials.2021.120776](https://doi.org/10.1016/j.biomaterials.2021.120776)

Provided by Pohang University of Science & Technology (POSTECH)

Citation: Engineering diseased human skin in vitro (2021, May 10) retrieved 9 April 2024 from <https://medicalxpress.com/news/2021-05-diseased-human-skin-vitro.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|