


Risk of second stroke can be reduced with prevention efforts based on cause of first stroke

May 24 2021



A blood clot forming in the carotid artery. Credit: American Heart Association

Having a stroke or a transient ischemic attack (TIA), sometimes called a "mini-stroke," increases the risk for a stroke in the future. Identifying

the cause of the stroke or TIA can lead to specific prevention strategies to reduce the risk of additional strokes, according to an updated guideline from the American Heart Association/American Stroke Association. The guideline is published today in *Stroke*.

Ischemic strokes account for 87% of strokes in the United States. An ischemic stroke occurs when [blood flow](#) in a vessel leading to the brain is blocked, by either clots or plaques. Strokes can lead to serious disability and/or death. A [transient ischemic attack](#), commonly referred to as a TIA, occurs when an artery is blocked for a short amount of time; thus, the blockage is transient (temporary) and does not cause permanent brain injury.

As prevention strategies have improved, studies have noted a reduction in recurrent stroke rates from 8.7% in the 1960s to 5.0% in the 2000s. Yet many [risk factors](#) for a second stroke remain poorly managed among stroke survivors.

A new recommendation of the "2021 Guideline for the Prevention of Stroke in Patients With Stroke and Transient Ischemic Attack" is for [health care professionals](#) to perform diagnostic evaluations to determine the cause of the first stroke or TIA within 48 hours of symptom onset. The guideline includes a section outlining treatment recommendations based on the cause of the initial stroke/TIA. Underlying causes could be related to blockages in large arteries in the neck or brain, small arteries in the brain damaged from [high blood pressure](#) or diabetes, irregular heart rhythms and many other potential causes.

"It is critically important to understand the best ways to prevent another stroke once someone has had a stroke or a TIA," said Dawn O. Kleindorfer, M.D., FAHA, chair of the guideline writing group, and professor and chair of the department of neurology at the University of Michigan School of Medicine in Ann Arbor, Michigan. "If we can

pinpoint the cause of the first stroke or TIA, we can tailor strategies to prevent a second stroke."

For patients who have survived a stroke or TIA, the secondary prevention guidelines recommend:

Managing their vascular risk factors, especially high blood pressure, as well as Type 2 diabetes, cholesterol, triglyceride levels and not smoking.

Limiting salt intake and/or following a Mediterranean diet.

If they are capable of [physical activity](#), engaging in moderate-intensity aerobic activity for at least 10 minutes four times a week or vigorous-intensity aerobic activity for at least 20 minutes twice a week.

"In fact, approximately 80% of strokes can be prevented by controlling blood pressure, eating a healthy diet, engaging in regular physical activity, not smoking and maintaining a healthy weight," said Amytis Towfighi, M.D., FAHA, vice-chair of the guideline writing group, and director of neurological services at the Los Angeles County Department of Health Services.

For health care professionals, the updated treatment recommendations highlighted in the guideline include:

Using multidisciplinary care teams to personalize care for patients and employing shared decision-making with the patient to develop care plans that incorporate a patient's wishes, goals and concerns.

Screening for and diagnosing atrial fibrillation (an irregular heart rhythm) and starting blood-thinning medications to reduce recurrent events.

Prescribing antithrombotic therapy, including antiplatelet medications (blood thinners) or anticoagulant medications (to prevent blood clotting), for nearly all patients who don't have contraindications. However, the combination of antiplatelets and anticoagulation is typically not recommended for preventing second strokes, and dual antiplatelet therapy, taking aspirin along with a second medicine to prevent blood clotting, is recommended short-term, only for specific patients: those with early arriving minor stroke and high-risk TIA or severe symptomatic stenosis.

Carotid endarterectomy, surgical removal of a blockage or, in select cases, the use of a stent in the carotid artery, should be considered for patients with narrowing arteries in the neck.

Aggressive medical management of risk factors and short-term dual antiplatelet therapy are preferred for patients with severe intracranial stenosis thought to cause the stroke or TIA.

In some patients, it is now reasonable to consider percutaneously closing (a less invasive, catheter-based surgical procedure) a patent foramen ovale, a small and fairly common heart defect.

The guideline is accompanied by a systematic review article, published simultaneously, "Benefits and Risks of Dual Versus Single Antiplatelet Therapy for Secondary Stroke Prevention." The review paper, chaired, by Devin L. Brown, M.D., M.S., is a meta-analysis of three short-duration clinical trials on dual antiplatelet therapy (DAPT) and concludes DAPT may be appropriate for select patients. The review authors note: "Additional research is needed to determine: the optimal timing of starting treatment relative to the clinical event; the optimal duration of DAPT to maximize the risk-benefit ratio; whether additional populations excluded from POINT and CHANCE [two of the trials examined], such as those with major stroke, may also benefit from early

DAPT; and whether certain genetic profiles eliminate the benefit of early DAPT."

"The secondary prevention of [stroke](#) guideline is one of the American Stroke Association's 'flagship' guidelines, last updated in 2014," Kleindorfer added. "There are also a number of changes to the writing and formatting of this guideline to make it easier for professionals to understand and locate information more quickly, ultimately greatly improving patient care and preventing more strokes in our patients."

More information: *Stroke* (2021). [DOI: 10.1161/STR.0000000000000375](#)

Provided by American Heart Association

Citation: Risk of second stroke can be reduced with prevention efforts based on cause of first stroke (2021, May 24) retrieved 24 June 2024 from <https://medicalxpress.com/news/2021-05-efforts-based.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--