


All in favor of a test for aortic aneurysms, raise your hand

May 26 2021, by Jim Shelton



An illustration of an aortic aneurysm and a demonstration of the thumb-palm test. Credit: Yale University

A Yale study has found that a self-conducted test involving the thumb and palm of one hand could help alert thousands of people each year to their risk of having a hidden aortic aneurysm.

Aortic aneurysms are the 13th most common cause of death among Americans, killing about 10,000 people each year. They affect adults of all ages.

If an aneurysm is detected early enough, a patient can be kept safe via restricting exercise, radiographic monitoring, or surgery.

But aortic aneurysms are hard to detect in advance, physicians say.

"The biggest problem in aneurysm disease is recognizing affected individuals within the [general population](#) before the aneurysm ruptures," said Dr. John A. Elefteriades, the William W.L. Glenn Professor of Surgery at Yale and emeritus director of the Aortic Institute at Yale New Haven Hospital. He is senior author of the new study, published in the *American Journal of Cardiology*.

For at least the past two decades, Elefteriades said, he and his colleagues have included the thumb-palm [test](#) in lectures to [medical students](#) and used the test for patients who might be at risk of carrying an aneurysm.

Yet despite its frequent usage, the test's accuracy had not been evaluated in a [clinical setting](#), the researchers said.

The test is simple: Holding up one hand and keeping the palm flat, the patient flexes their thumb as far as possible across the palm. If the thumb crosses beyond the far edge of the flat palm, the patient may be harboring a hidden aneurysm.

Being able to move the thumb in that way is an indication that a patient's long bones are excessive and their joints are lax—possible signs of connective tissue disease throughout the body, including the aorta.

The researchers said it is important to emphasize that not everyone who

tests positive is an aneurysm carrier. Also, they noted, aneurysms often take decades to progress to the point of rupture and a positive test is not cause for panic.

For the study, the researchers gave the test to 305 patients undergoing cardiac surgery for a variety of disorders, including [aortic aneurysms](#). "Our study showed that the majority of aneurysm patients do not manifest a positive thumb-palm sign, but patients who do have a positive test have a high likelihood of harboring an aneurysm," Elefteriades said.

Based on their findings, Elefteriades and his colleagues said the thumb-palm test should be included in standard physical examinations of patients—particularly those who have a family history of aortic aneurysm.

"Spreading knowledge of this test may well identify silent [aneurysm](#) carriers and save lives," Elefteriades said.

More information: Rachel Blumel et al, Accuracy of the "Thumb-Palm Test" for Detection of Ascending Aortic Aneurysm, *The American Journal of Cardiology* (2021). [DOI: 10.1016/j.amjcard.2021.03.041](https://doi.org/10.1016/j.amjcard.2021.03.041)

Provided by Yale University

Citation: All in favor of a test for aortic aneurysms, raise your hand (2021, May 26) retrieved 25 April 2024 from <https://medicalxpress.com/news/2021-05-favor-aortic-aneurysms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.