


Researcher identifies new mode of transmission for bacteria

May 21 2021



This scanning electron microscope image shows the characteristic spiral, or corkscrew, shape of *C. jejuni* cells and related structures. Credit: De Wood; digital colorization by Chris Pooley/ Public Domain

Campylobacter infection, one of the most common foodborne illnesses in the Western world, can also be spread through sexual contact, according to a new research discovery by an OU Hudson College of Public Health faculty member, working in conjunction with colleagues in Denmark.

The team's research has been published in *Emerging Infectious Diseases*, a journal published by the Centers for Disease Control and Prevention (CDC), and is the first known study to prove this mode of transmission for Campylobacter. During a time when COVID-19 has dominated news about infectious diseases, the research is a reminder that many other pathogens affect lives around the world every day. The study was led by infectious disease epidemiologist Katrin Kuhn, Ph.D., an assistant professor in the Department of Biostatistics and Epidemiology in the OU Hudson College of Public Health.

"This research is important for public health messaging and for physicians as they talk to their patients about risks associated with sexual contact," Kuhn said. "Although Campylobacter infection is usually not a serious disease, it causes diarrhea, which can result in people missing work, losing productivity or perhaps losing their job. It poses an additional risk for people with underlying health conditions."

Campylobacter infections usually occur when people eat chicken that has not been cooked thoroughly or when juices from uncooked poultry make their way into other food. Infections can also be caused by drinking unpasteurized milk or water that has been contaminated by the

feces of infected animals. However, those didn't account for all cases of infection, Kuhn said, and she wondered if there was another route of transmission that remained unproven. An outbreak of *Campylobacter* infections in northern Europe among men who have sex with men prompted her to study that population of people in Denmark, where she was working when the research began.

The study results showed that the rate of *Campylobacter* infection was 14 times higher in men who have sex with men than the control subjects. Although the study focused on men who have sex with men, the results are relevant to people of any sexual orientation who engage in sexual behavior that may involve fecal-oral contact, Kuhn said.

Two other bacteria, *Salmonella* and *Shigella*, were used as comparisons in the study. *Salmonella* is spread primarily through infected foods, while *Shigella* can be transmitted through food or sexual contact. *Salmonella* has a high infectious dose, meaning people must ingest a significant amount of the bacteria before they become ill. However, *Shigella* and *Campylobacter* have low infectious doses, which makes transmission easier.

"That's an additional reason why we believe *Campylobacter* can be transmitted through [sexual contact](#) like *Shigella* is—because people can become infected when only small amounts of the bacteria are present," Kuhn said.

Campylobacter infections are probably more prevalent than the numbers show. For every one person who goes to the doctor and is diagnosed, epidemiologists estimate that 20 more people are infected, Kuhn said. Although treatment is usually required only for severe cases, complications can occur, especially in people who have compromised immune systems. In some cases, [infection](#) can result in reactive arthritis, in which the body's immune system attacks itself, causing pain in the

joints. Infection can also lead to Guillain-Barré Syndrome, a serious nerve disorder that can cause paralysis.

"This is an interesting time because COVID-19 has made people more aware of the importance of monitoring infectious diseases in general, not only during a pandemic," she said. "There are many infections like the one caused by *Campylobacter* that make people sick. It's important that we spotlight the fact that these diseases exist and that we continue to conduct research on their effects and modes of transmission."

Before arriving at the OU Hudson College of Public Health, Kuhn served as a senior infectious [disease](#) epidemiologist at Statens Serum Institut in Denmark. Her work focused on food- and water-borne infections, and she was responsible for the national surveillance of *Campylobacter* and *Shigella*. She began this study while in Denmark and completed it after moving to Oklahoma. Statens Serum Institut is the Danish national institute for infectious diseases and the primary institute for surveillance of and research on infectious diseases in Denmark.

"A formal collaboration between OU Hudson College of Public Health and Statens Serum Institut will build a solid foundation for strengthening transatlantic research and, not least, improving the way that we monitor, understand and prevent [infectious diseases](#) in Oklahoma," Kuhn said.

More information: Katrin Gaardbo Kuhn et al, Sexual Contact as Risk Factor for *Campylobacter* Infection, Denmark, *Emerging Infectious Diseases* (2021). [DOI: 10.3201/eid2704.202337](https://doi.org/10.3201/eid2704.202337)

Provided by University of Oklahoma

Citation: Researcher identifies new mode of transmission for bacteria (2021, May 21) retrieved

26 April 2024 from <https://medicalxpress.com/news/2021-05-mode-transmission-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.