

Low levels of omega-3 associated with higher risk of psychosis, says study

May 31 2021



Credit: CC0 Public Domain

New research has found that adolescents with higher levels of an omega-3 fatty acid in their blood were less likely to develop psychotic disorder in early adulthood, suggesting that it may have a potential

preventative effect of reducing the risk of psychosis.

The study, led by researchers from RCSI University of Medicine and Health Sciences, is published in *Translational Psychiatry*.

Over 3,800 individuals in Bristol's Children of the 90s health study were assessed for [psychotic disorder](#), [depressive disorder](#) and generalized anxiety disorder at age 17 and at age 24.

During these assessments, blood samples were collected, and the researchers measured the levels of omega-6 fatty acids, which generally increase inflammation in the body, and omega-3 fatty acids, which generally reduce inflammation.

While there was little evidence that fatty acids were associated with mental disorders at age 17, the researchers found that 24-year-olds with psychotic disorder, depressive disorder and generalized anxiety disorder had higher levels of omega-6 than omega-3 fatty acids compared to those without these disorders.

The researchers also found that 24-year-olds with psychotic disorder had lower levels of DHA, an omega-3 fatty [acid](#) typically found in oily fish or dietary supplements, than 24-year-olds without psychotic disorder. In a group of over 2,700 individuals who were tracked over time, adolescents with higher levels of DHA at age 17 were 56% less likely to develop psychotic disorder seven years later at age 24. This suggests that DHA in adolescence may have a potential preventative effect of reducing the risk of psychosis in [early adulthood](#).

These results remained consistent when accounting for other factors such as sex, body mass index, tobacco smoking and socio-economic status.

"The study needs to be replicated, but if the findings are consistent, these results would suggest that enhanced dietary intake of omega-3 fatty acids among adolescents, such as through oily fish like mackerel, could prevent some people from developing psychosis in their early twenties," said Professor David Cotter, senior author of the study and professor molecular psychiatry at RCSI.

"The results could also raise questions about the relationship between the development of mental health [disorders](#) and omega-6 fatty acids, which are typically found in vegetable oils."

David Mongan, RCSI Ph.D. student and Irish Clinical Academic Training (ICAT) Fellow, analyzed the data with the supervision of Professor David Cotter and Professor Mary Cannon from the RCSI Department of Psychiatry. The ICAT program is supported by the Wellcome Trust and the Health Research Board, the Health Service Executive National Doctors Training and Planning and the Health and Social Care, Research and Development Division, Northern Ireland.

"We need to do more research to learn about the mechanisms behind this effect, but it could possibly be related to reducing inflammation or decreasing inappropriate pruning of brain connections during adolescence," said Dr. David Mongan, the study's first author, who is a psychiatry trainee and Ph.D. student at RCSI.

More information: David Mongan et al, Plasma polyunsaturated fatty acids and mental disorders in adolescence and early adulthood: cross-sectional and longitudinal associations in a general population cohort, *Translational Psychiatry* (2021). [DOI: 10.1038/s41398-021-01425-4](https://doi.org/10.1038/s41398-021-01425-4)

Provided by RCSI University of Medicine and Health Sciences

Citation: Low levels of omega-3 associated with higher risk of psychosis, says study (2021, May 31) retrieved 24 April 2024 from <https://medicalxpress.com/news/2021-05-omega-higher-psychosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.