

Long term use of prescription meds for insomnia not linked to better quality sleep

May 12 2021



Credit: CC0 Public Domain

Long term use of prescription meds for insomnia doesn't seem to improve disturbed sleep in middle-aged women, suggests research published in the online journal *BMJ Open*.

There was no difference in [sleep quality](#) or duration between those who did and didn't take these meds for 1 to 2 years, the findings show.

Disturbed sleep—difficulty falling and/or staying asleep and waking early—is common. An estimated 9 million adults in the US alone say they take prescription meds to try and get a good night's sleep.

Poor quality sleep is associated with ill health, including diabetes, [high blood pressure](#), pain and depression, and various drugs are prescribed to induce shut-eye.

These include benzodiazepines, Z-drugs which include zolpidem, zaleplon and eszopiclone, as well as other agents mostly intended for other conditions (off label use), such as quelling anxiety and depression.

The clinical trial data indicate that many of these drugs work in the short term (up to 6 months), but insomnia can be chronic, and many people take these drugs for longer, say the researchers.

They therefore wanted to assess the effectiveness of drugs used to tackle insomnia over the long term among an ethnically diverse group of middle aged [women](#) who developed [sleep disorders](#).

The women were all part of the Study of Women's Health Across the Nation (SWAN), a long term multicentre study looking at biological and psychosocial changes arising during the menopause. The women's average age was 49.5 and around half were white.

Sleep disturbances were defined as difficulty falling asleep, frequent awakening, and waking up early and rated on a 5-point scale, ranging from no difficulty on any night (1) to difficulty on 5 or more nights of the week (5), reported during an average of 21 years of monitoring.

Sleep disturbances, as measured on the ratings scale, were compared among those who did and didn't take prescription meds to improve their sleep after 1 and 2 years.

Some 238 women who started using medication to tackle insomnia during the monitoring period were matched with 447 women who didn't take these drugs.

Both groups of women reported difficulty falling asleep on 1 out of every 3 nights, waking frequently on 2 out of 3 nights, and waking up early on 1 in every 3 nights of the week. More than 70% of women in both groups reported [disturbed sleep](#) at least 3 times a week.

To begin with, sleep disturbance ratings were similar between the two groups of women. Those taking prescription meds for their sleep problems had average scores for difficulty falling asleep, waking up frequently, and for waking up early of 2.7, 3.8, and 2.8 respectively.

This compares with equivalent ratings of 2.6, 3.7, and 2.7, respectively, for those not taking prescription meds to get a good night's sleep.

After 1 year, average ratings among those taking the meds were 2.6, 3.6, and 2.8, respectively. The equivalent average scores among those not using prescription meds for their sleep problems were 2.3, 3.5, and 2.5, respectively.

None of the 1 year changes was statistically significant nor did they differ between the two groups. And after 2 years there were no statistically significant reductions in sleep disturbances among those taking prescription meds compared with those who didn't.

This is an observational study, and as such can't establish cause, only correlation. What's more, around half of the women were current or

former smokers and 1 in 5 were moderate to heavy drinkers, both of which may affect sleep quality.

Information collected on prescription meds was also collected only at annual or biennial study visits, and there may have been intermittent or periods of no use between visits, say the researchers. Nor were there any objective measures of sleep quality.

Nevertheless, conclude the researchers: "Sleep disturbances are common and increasing in prevalence. The use of sleep medications has grown, and they are often used over a long period, despite the relative lack of evidence from [randomised controlled trials]."

These drugs may work well in some people with sleep disturbances over several years, but the findings of this study should give pause for thought to prescribing clinicians and patients thinking about taking prescription meds for sleep disturbances in middle age, they add.

More information: Daniel H Solomon et al. Prescription medications for sleep disturbances among midlife women during 2 years of follow-up: a SWAN retrospective cohort study, *BMJ Open* (2021). [DOI: 10.1136/bmjopen-2020-045074](https://doi.org/10.1136/bmjopen-2020-045074)

Provided by British Medical Journal

Citation: Long term use of prescription meds for insomnia not linked to better quality sleep (2021, May 12) retrieved 7 May 2024 from <https://medicalxpress.com/news/2021-05-term-prescription-meds-insomnia-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.